

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
18 November 2004 (18.11.2004)

PCT

(10) International Publication Number
WO 2004/098507 A2

- (51) International Patent Classification⁷: A61K
- (21) International Application Number:
PCT/US2004/013345
- (22) International Filing Date: 30 April 2004 (30.04.2004)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/466,641 30 April 2003 (30.04.2003) US
- (71) Applicant (for all designated States except US): BIOX-
ELL S.P.A. [IT/IT]; Via Olgettina, 58, I-20132 Milano
(IT).
- (72) Inventors; and
- (73) Inventors/Applicants (for US only): ADORINI, Lu-
ciano [IT/IT]; via Augusto Righi, 8, I-20133 Milano (IT).
PENNA, Giuseppe [IT/IT]; Via Cervino 23, I-20095
Milano (IT). USKOKOVIC, Milan, R. [US/US]; 253
Highland Avenue, Upper Montclair, NJ 07043. (US).
MAEHR, Hubert [US/US]; 30 Balsam Road, Wayne, NJ
07470 (US).
- (74) Agents: LAURO, Peter, C. et al.; Edwards & Angell,
L.L.P., P.O. Box 55874, Boston, MA 02205 (US).


- (81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EG, ES, FI,
GB, GD, GH, GM, GR, HU, ID, IL, IN, IS, JP, KE,
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,
MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG,
PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM,
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM,
ZW.
- (84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,
FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI,
SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished
upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: 1,3 ACYLATED 24-KETO-VITAMIN D₃ COMPOUNDS AND METHODS OF USE THEREOF



(I)

(57) Abstract: The invention provides 1,3 acylated, 24-keto vitamin D₃ compounds of formula I: wherein: X₁ and X₂ are each independently H₂ or =CH₂, provided X₁ and X₂ are not both =CH₂; R₁ and R₂ are each independently, hydroxyl, OC(O)C₁-C₄ alkyl, OC(O)hydroxyalkyl, OC(O)haloalkyl, provided that R₁ and R₂ are not both hydroxyl; R₃ and R₄ are each independently hydrogen, C₁-C₄ alkyl, hydroxyalkyl, or haloalkyl, or R₃ and R₄ taken together with C₂₀ form C₃-C₆ cycloalkyl; and R₅ and R₆ are each independently C₁-C₄ alkyl, hydroxyalkyl, or haloalkyl; and pharmaceutically acceptable esters, salts, and prodrugs thereof. Methods for using the compounds to treat vitamin D₃ associated states, and pharmaceutical compositions

containing the compounds are also disclosed.

1,3 ACETYLATED, 24-KETO-VITAMIN D₃ COMPOUNDS AND METHODS OF USE THEREOF

Related Application

5 This application claims the benefit of U.S. provisional patent application Ser. No. 60/466,641 filed April 30, 2003, the disclosure of which application is incorporated herein in its entirety by this reference.

Background of the Invention

10 The importance of vitamin D (cholecalciferol) in the biological systems of higher animals has been recognized since its discovery by Mellanby in 1920 (Mellanby, E. (1921) *Spec. Rep. Ser. Med. Res. Council* (GB) SRS 61:4). It was in the interval of 1920-1930 that vitamin D officially became classified as a "vitamin" that was essential for the normal development of the skeleton and maintenance of calcium and
15 phosphorous homeostasis.

 Studies involving the metabolism of vitamin D₃ were initiated with the discovery and chemical characterization of the plasma metabolite, 25-hydroxyvitamin D₃ [25(OH)D₃] (Blunt, J.W. *et al.* (1968) *Biochemistry* 6:3317-3322) and the hormonally active form, 1 α ,25(OH)₂D₃ (Myrtle, J.F. *et al.* (1970) *J. Biol. Chem.* 245:1190-1196;
20 Norman, A.W. *et al.* (1971) *Science* 173:51-54; Lawson, D.E.M. *et al.* (1971) *Nature* 230:228-230; Holick, M.F. (1971) *Proc. Natl. Acad. Sci. USA* 68:803-804). The formulation of the concept of a vitamin D endocrine system was dependent both upon appreciation of the key role of the kidney in producing 1 α , 25(OH)₂D₃ in a carefully regulated fashion (Fraser, D.R. and Kodicek, E (1970) *Nature* 288:764-766; Wong, R.G.
25 *et al.* (1972) *J. Clin. Invest.* 51:1287-1291), and the discovery of a nuclear receptor for 1 α ,25(OH)₂D₃ (VD₃R) in the intestine (Haussler, M.R. *et al.* (1969) *Exp. Cell Res.* 58:234-242; Tsai, H.C. and Norman, A.W. (1972) *J. Biol. Chem.* 248:5967-5975).

 The operation of the vitamin D endocrine system depends on the following: first, on the presence of cytochrome P450 enzymes in the liver (Bergman, T. and Postlind, H.
30 (1991) *Biochem. J.* 276:427-432; Ohyama, Y and Okuda, K. (1991) *J. Biol. Chem.* 266:8690-8695) and kidney (Henry, H.L. and Norman, A.W. (1974) *J. Biol. Chem.* 249:7529-7535; Gray, R.W. and Ghazarian, J.G. (1989) *Biochem. J.* 259:561-568), and

in a variety of other tissues to effect the conversion of vitamin D₃ into biologically active metabolites such as 1 α , 25(OH)₂D₃ and 24R,25(OH)₂D₃; second, on the existence of the plasma vitamin D binding protein (DBP) to effect the selective transport and delivery of these hydrophobic molecules to the various tissue components of the vitamin D endocrine system (Van Baelen, H. *et al.* (1988) *Ann NY Acad. Sci.* 538:60-68; Cooke, N.E. and Haddad, J.G. (1989) *Endocr. Rev.* 10:294-307; Bikle, D.D. *et al.* (1986) *J. Clin. Endocrinol. Metab.* 63:954-959); and third, upon the existence of stereoselective receptors in a wide variety of target tissues that interact with the agonist 1 α ,25(OH)₂D₃ to generate the requisite specific biological responses for this secosteroid hormone (Pike, J.W. (1991) *Annu. Rev. Nutr.* 11:189-216). To date, there is evidence that nuclear receptors for 1 α ,25(OH)₂D₃ (VD₃R) exist in more than 30 tissues and cancer cell lines (Reichel, H. and Norman, A.W. (1989) *Annu. Rev. Med.* 40:71-78).

Vitamin D₃ and its hormonally active forms are well-known regulators of calcium and phosphorous homeostasis. These compounds are known to stimulate, at least one of, intestinal absorption of calcium and phosphate, mobilization of bone mineral, and retention of calcium in the kidneys. Furthermore, the discovery of the presence of specific vitamin D receptors in more than 30 tissues has led to the identification of vitamin D₃ as a pluripotent regulator outside its classical role in calcium/bone homeostasis. A paracrine role for 1 α ,25(OH)₂D₃ has been suggested by the combined presence of enzymes capable of oxidizing vitamin D₃ into its active forms, *e.g.*, 25-OHD-1 α -hydroxylase, and specific receptors in several tissues such as bone, keratinocytes, placenta, and immune cells. Moreover, vitamin D₃ hormone and active metabolites have been found to be capable of regulating cell proliferation and differentiation of both normal and malignant cells (Reichel, H. *et al.* (1989) *Ann. Rev. Med.* 40: 71-78).


Given the activities of vitamin D₃ and its metabolites, much attention has focused on the development of synthetic analogs of these compounds. A large number of these analogs involve structural modifications in the A ring, B ring, C/D rings, and, primarily, the side chain (Bouillon, R. *et al.*, *Endocrine Reviews* 16(2):201-204). Although a vast majority of the vitamin D₃ analogs developed to date involve structural modifications in the side chain, a few studies have reported the biological profile of A-ring diastereomers (Norman, A.W. *et al.* *J. Biol. Chem.* 268 (27): 20022-20030).

Furthermore, biological esterification of steroids has been studied (Hochberg, R.B., (1998) *Endocr Rev.* 19(3): 331-348), and esters of vitamin D₃ are known (WO 97/11053).

Moreover, despite much effort in developing synthetic analogs, clinical applications of vitamin D and its structural analogs have been limited by the undesired side effects elicited by these compounds after administration to a subject for known indications/applications of vitamin D compounds.

Summary of the Invention

In one aspect, the invention provides a vitamin D₃ compound of the formula I:



wherein:

X₁ and X₂ are each independently H₂ or =CH₂, provided X₁ and X₂ are not both =CH₂;

R₁ and R₂ are each independently, hydroxyl, OC(O)C₁-C₄ alkyl, OC(O)hydroxyalkyl, OC(O)haloalkyl, provided that R₁ and R₂ are not both hydroxyl;

R₃ and R₄ are each independently hydrogen, C₁-C₄ alkyl, hydroxyalkyl, or haloalkyl, or R₃ and R₄ taken together with C₂₀ form C₃-C₆ cycloalkyl; and

R₅ and R₆ are each independently C₁-C₄ alkyl, hydroxyalkyl, or haloalkyl; and pharmaceutically acceptable esters, salts, and prodrugs thereof.

In another aspect, the invention provides a method for treating a subject for a vitamin D₃ associated state. The method includes administering to the subject an

effective amount of a vitamin D₃ compound of formula I, such that the subject is treated for the vitamin D₃ associated state.

In yet another aspect, the invention provides a pharmaceutical composition. The composition comprises an effective amount of a vitamin D₃ compound of formula I, and
5 a pharmaceutically acceptable carrier.

In a further aspect, the method provides a method of ameliorating a deregulation of calcium and phosphate metabolism. The method includes administering to a subject a therapeutically effective amount of a vitamin D₃ compound of formula I, so as to ameliorate the deregulation of the calcium and phosphate metabolism.

10 In another aspect, the invention provides a method of modulating the expression of an immunoglobulin-like transcript 3 (ILT3) surface molecule in a cell. The method includes contacting the cell with a vitamin D₃ compound of formula I in an amount effective to modulate the expression of an immunoglobulin-like transcript 3 (ILT3) surface molecule in the cell.

15 In yet another aspect, the invention provides a method of treating an ILT3-associated disorder in a subject. The method includes administering to the subject a vitamin D₃ compound of formula I in an amount effective to modulate the expression of an ILT3 surface molecule, thereby treating the ILT3-associated disorder in the subject.

In still another aspect, the invention provides a method of inducing
20 immunological tolerance in a subject. The method includes administering to the subject a vitamin D₃ compound of formula I in an amount effective to modulate the expression of an ILT3 surface molecule, thereby inducing immunological tolerance in the subject.

In a further aspect, the invention provides a method of inhibiting transplant rejection in a subject. The method includes administering to the subject a vitamin D₃
25 compound of formula I in an amount effective to modulate the expression of an ILT3 surface molecule, thereby inhibiting transplant rejection in the subject.

In yet another aspect, the invention provides a packaged formulation for use in the treatment of a vitamin D₃ associated state. The packaged formulation includes a pharmaceutical composition comprising a vitamin D₃ compound of formula I and a
30 pharmaceutically-acceptable carrier, packaged with instructions for use in the treatment of a vitamin D₃ associated state.

In another aspect, the invention provides a packaged formulation for use in the treatment of an ILT-3 associated disorder. The packed formulation includes a

pharmaceutical composition comprising a vitamin D₃ compound of formula I and a pharmaceutically-acceptable carrier, packaged with instructions for use in the treatment of an ILT3-associated disorder.

- In a further aspect, the invention provides a method for modulating immunosuppressive activity by an antigen-presenting cell. The method includes contacting an antigen-presenting cell with a vitamin D₃ compound of formula I in an amount effective to modulate ILT3 surface molecule expression, thereby modulating the immunosuppressive activity by the antigen-presenting cell.

10 Detailed Description of the Invention

I. DEFINITIONS

- Before further description of the present invention, and in order that the invention may be more readily understood, certain terms are first defined and collected here for convenience.

- The term "administration" or "administering" includes routes of introducing the vitamin D₃ compound(s) to a subject to perform their intended function. Examples of routes of administration which can be used include injection (subcutaneous, intravenous, parenterally, intraperitoneally, intrathecal), oral, inhalation, rectal and transdermal. The pharmaceutical preparations are, of course, given by forms suitable for each administration route. For example, these preparations are administered in tablets or capsule form, by injection, inhalation, eye lotion, ointment, suppository, etc. administration by injection, infusion or inhalation; topical by lotion or ointment; and rectal by suppositories. Oral administration is preferred. The injection can be bolus or can be continuous infusion. Depending on the route of administration, the vitamin D₃ compound can be coated with or disposed in a selected material to protect it from natural conditions which may detrimentally effect its ability to perform its intended function. The vitamin D₃ compound can be administered alone, or in conjunction with either another agent as described above or with a pharmaceutically-acceptable carrier, or both. The vitamin D₃ compound can be administered prior to the administration of the other agent, simultaneously with the agent, or after the administration of the agent. Furthermore, the vitamin D₃ compound can also be administered in a proform which is converted into its active metabolite, or more active metabolite *in vivo*.

The term "alkyl" refers to the radical of saturated aliphatic groups, including straight-chain alkyl groups, branched-chain alkyl groups, cycloalkyl (alicyclic) groups, alkyl substituted cycloalkyl groups, and cycloalkyl substituted alkyl groups. The term alkyl further includes alkyl groups, which can further include oxygen, nitrogen, sulfur or phosphorous atoms replacing one or more carbons of the hydrocarbon backbone, *e.g.*, oxygen, nitrogen, sulfur or phosphorous atoms. In preferred embodiments, a straight chain or branched chain alkyl has 30 or fewer carbon atoms in its backbone (*e.g.*, C₁-C₃₀ for straight chain, C₃-C₃₀ for branched chain), preferably 26 or fewer, and more preferably 20 or fewer. Likewise, preferred cycloalkyls have from 3-10 carbon atoms in their ring structure, and more preferably have 3, 4, 5, 6 or 7 carbons in the ring structure.

Moreover, the term alkyl as used throughout the specification and claims is intended to include both "unsubstituted alkyls" and "substituted alkyls," the latter of which refers to alkyl moieties having substituents replacing a hydrogen on one or more carbons of the hydrocarbon backbone. Such substituents can include, for example, halogen, hydroxyl, alkylcarbonyloxy, arylcarbonyloxy, alkoxycarbonyloxy, aryloxy carbonyloxy, carboxylate, alkylcarbonyl, alkoxycarbonyl, aminocarbonyl, alkylthiocarbonyl, alkoxyl, phosphate, phosphonato, phosphinato, cyano, amino (including alkyl amino, dialkylamino, arylamino, diarylamino, and alkylarylamino), acylamino (including alkylcarbonylamino, arylcarbonylamino, carbamoyl and ureido), amidino, imino, sulfhydryl, alkylthio, arylthio, thiocarboxylate, sulfates, sulfonato, sulfamoyl, sulfonamido, nitro, trifluoromethyl, cyano, azido, heterocyclyl, alkylaryl, or an aromatic or heteroaromatic moiety. It will be understood by those skilled in the art that the moieties substituted on the hydrocarbon chain can themselves be substituted, if appropriate. Cycloalkyls can be further substituted, *e.g.*, with the substituents described above. An "alkylaryl" moiety is an alkyl substituted with an aryl (*e.g.*, phenylmethyl (benzyl)). The term "alkyl" also includes unsaturated aliphatic groups analogous in length and possible substitution to the alkyls described above, but that contain at least one double or triple bond respectively.

Unless the number of carbons is otherwise specified, "lower alkyl" as used herein means an alkyl group, as defined above, but having from one to ten carbons, more preferably from one to six, and most preferably from one to four carbon atoms in its backbone structure, which may be straight or branched-chain. Examples of lower alkyl groups include methyl, ethyl, n-propyl, i-propyl, tert-butyl, hexyl, heptyl, octyl and

so forth. In preferred embodiment, the term "lower alkyl" includes a straight chain alkyl having 4 or fewer carbon atoms in its backbone, *e.g.*, C₁-C₄ alkyl.

The terms "alkoxyalkyl," "polyaminoalkyl" and "thioalkoxyalkyl" refer to alkyl groups, as described above, which further include oxygen, nitrogen or sulfur atoms replacing one or more carbons of the hydrocarbon backbone, *e.g.*, oxygen, nitrogen or sulfur atoms.

The terms "alkenyl" and "alkynyl" refer to unsaturated aliphatic groups analogous in length and possible substitution to the alkyls described above, but that contain at least one double or triple bond, respectively. For example, the invention contemplates cyano and propargyl groups.

The term "antigen" includes a substance which elicits an immune response. The antigens of the invention to which tolerance is induced may or may not be exogenously derived relative to the host. For example, the method of the invention may be used to induce tolerance to an "autoantigen." An autoantigen is a normal constituent of the body that reacts with an autoantibody. The invention also includes inducing tolerance to an "alloantigen." Alloantigen refers to an antigen found only in some members of a species, for example the blood group substances. An allograft is a graft to a genetically different member of the same species. Allografts are rejected by virtue of the immunological response of T lymphocytes to histocompatibility antigens. The method of the invention also provides for inducing tolerance to a "xenoantigen." Xenoantigens are substances that cause an immune reaction due to differences between different species. Thus, a xenograft is a graft from a member of one species to a member of a different species. Xenografts are usually rejected within a few days by antibodies and cytotoxic T lymphocytes to histocompatibility antigens.

The language "antigen-presenting cell" or "APC" includes a cell that is able to present an antigen to, for example, a T helper cell. Antigen-presenting cells include B lymphocytes, accessory cells or non-lymphocytic cells, such as dendritic cells, Langerhans cells, and mononuclear phagocytes that help in the induction of an immune response by presenting antigen to helper T lymphocytes. The antigen-presenting cell of the present invention is preferably of myeloid origin, and includes, but is not limited to, dendritic cells, macrophages, monocytes. APCs of the present invention may be isolated from the bone marrow, blood, thymus, epidermis, liver, fetal liver, or the spleen.

The terms "antineoplastic agent" and "antiproliferative agent" are used interchangeably herein and includes agents that have the functional property of inhibiting the proliferation of a vitamin D₃-responsive cells, e.g., inhibit the development or progression of a neoplasm having such a characteristic, particularly a hematopoietic neoplasm.

The term "aryl" as used herein, refers to the radical of aryl groups, including 5- and 6-membered single-ring aromatic groups that may include from zero to four heteroatoms, for example, benzene, pyrrole, furan, thiophene, imidazole, benzoxazole, benzothiazole, triazole, tetrazole, pyrazole, pyridine, pyrazine, pyridazine and pyrimidine, and the like. Aryl groups also include polycyclic fused aromatic groups such as naphthyl, quinolyl, indolyl, and the like. Those aryl groups having heteroatoms in the ring structure may also be referred to as "aryl heterocycles," "heteroaryls" or "heteroaromatics." The aromatic ring can be substituted at one or more ring positions with such substituents as described above, as for example, halogen, hydroxyl, alkoxy, alkylcarbonyloxy, arylcarbonyloxy, alkoxycarbonyloxy, aryloxy, carboxylate, alkylcarbonyl, alkoxycarbonyl, aminocarbonyl, alkylthiocarbonyl, phosphate, phosphonato, phosphinato, cyano, amino (including alkyl amino, dialkylamino, arylamino, diarylamino, and alkylarylamino), acylamino (including alkylcarbonylamino, arylcarbonylamino, carbamoyl and ureido), amidino, imino, sulfhydryl, alkylthio, arylthio, thiocarboxylate, sulfates, sulfonato, sulfamoyl, sulfonamido, nitro, trifluoromethyl, cyano, azido, heterocyclyl, alkylaryl, or an aromatic or heteroaromatic moiety. Aryl groups can also be fused or bridged with alicyclic or heterocyclic rings which are not aromatic so as to form a polycycle (e.g., tetralin).

The language "autoimmune disease" or "autoimmune disorder" refers to the condition where the immune system attacks the host's own tissue(s). In an autoimmune disease, the immune tolerance system of the patient fails to recognize self antigens and, as a consequence of this loss of tolerance, brings the force of the immune system to bear on tissues which express the antigen. Autoimmune disorders include, but are not limited to, type 1 insulin-dependent diabetes mellitus, adult respiratory distress syndrome, inflammatory bowel disease, dermatitis, meningitis, thrombotic thrombocytopenic purpura, Sjogren's syndrome, encephalitis, uveitis, uveoretinitis, leukocyte adhesion deficiency, rheumatoid arthritis, rheumatic fever, Reiter's syndrome, psoriatic arthritis, progressive systemic sclerosis, primary biliary cirrhosis, pemphigus, pemphigoid,

necrotizing vasculitis, myasthenia gravis, multiple sclerosis, lupus erythematosus, polymyositis, sarcoidosis, granulomatosis, vasculitis, pernicious anemia, CNS inflammatory disorder, antigen-antibody complex mediated diseases, autoimmune haemolytic anemia, Hashimoto's thyroiditis, Graves disease, habitual spontaneous abortions, Reynard's syndrome, glomerulonephritis, dermatomyositis, chronic active hepatitis, celiac disease, autoimmune complications of AIDS, atrophic gastritis, ankylosing spondylitis and Addison's disease.

The language "biological activities" of vitamin D₃ includes all activities elicited by vitamin D₃ compounds in a responsive cell. It includes genomic and non-genomic activities elicited by these compounds (Gniadecki R. and Calverley M.J. (1998) *Pharmacology & Toxicology* 82: 173-176; Bouillon, R. *et al.* (1995) *Endocrinology Reviews* 16(2):206-207; Norman A.W. *et al.* (1992) *J. Steroid Biochem Mol. Biol* 41:231-240; Baran D.T. *et al.* (1991) *J. Bone Miner Res.* 6:1269-1275; Caffrey J.M. and Parach-Carson M.C. (1989) *J. Biol. Chem.* 264:20265-20274; Nemere I. *et al.* (1984) *Endocrinology* 115:1476-1483).

The language "bone metabolism" includes direct or indirect effects in the formation or degeneration of bone structures, *e.g.*, bone formation, bone resorption, *etc.*, which may ultimately affect the concentrations in serum of calcium and phosphate. This term is also intended to include effects of compounds of the invention in bone cells, *e.g.*, osteoclasts and osteoblasts, that may in turn result in bone formation and degeneration.

The language "calcium and phosphate homeostasis" refers to the careful balance of calcium and phosphate concentrations, intracellularly and extracellularly, triggered by fluctuations in the calcium and phosphate concentration in a cell, a tissue, an organ or a system. Fluctuations in calcium levels that result from direct or indirect responses to compounds of the invention are intended to be included by these terms.

The term "carcinoma" is art recognized and refers to malignancies of epithelial or endocrine tissues including respiratory system carcinomas, gastrointestinal system carcinomas, genitourinary system carcinomas, testicular carcinomas, breast carcinomas, prostatic carcinomas, endocrine system carcinomas, and melanomas. Exemplary carcinomas include those forming from tissue of the cervix, lung, prostate, breast, head and neck, colon and ovary. The term also includes carcinosarcomas, *e.g.*, which include malignant tumors composed of carcinomatous and sarcomatous tissues. An

"adenocarcinoma" refers to a carcinoma derived from glandular tissue or in which the tumor cells form recognizable glandular structures.

The term "chiral" refers to molecules which have the property of non-superimposability of the mirror image partner, while the term "achiral" refers to
5 molecules which are superimposable on their mirror image partner.

The term "diastereomers" refers to stereoisomers with two or more centers of dissymmetry and whose molecules are not mirror images of one another.

The term "effective amount" includes an amount effective, at dosages and for periods of time necessary, to achieve the desired result, *e.g.*, sufficient treat a vitamin D₃
10 associated state or to modulate ILT3 expression in a cell. An effective amount of vitamin D₃ compound may vary according to factors such as the disease state, age, and weight of the subject, and the ability of the vitamin D₃ compound to elicit a desired response in the subject. Dosage regimens may be adjusted to provide the optimum therapeutic response. An effective amount is also one in which any toxic or detrimental
15 effects (*e.g.*, side effects) of the angiogenesis inhibitor compound are outweighed by the therapeutically beneficial effects.

A therapeutically effective amount of vitamin D₃ compound (*i.e.*, an effective dosage) may range from about 0.001 to 30 µg/kg body weight, preferably about 0.01 to 25 µg/kg body weight, more preferably about 0.1 to 20 µg/kg body weight, and even
20 more preferably about 1 to 10 µg/kg, 2 to 9 µg/kg, 3 to 8 µg/kg, 4 to 7 µg/kg, or 5 to 6 µg/kg body weight. The skilled artisan will appreciate that certain factors may influence the dosage required to effectively treat a subject, including but not limited to the severity of the disease or disorder, previous treatments, the general health and/or age of the subject, and other diseases present. Moreover, treatment of a subject with a
25 therapeutically effective amount of a vitamin D₃ compound can include a single treatment or, preferably, can include a series of treatments. In one example, a subject is treated with a vitamin D₃ compound in the range of between about 0.1 to 20 µg/kg body weight, one time per week for between about 1 to 10 weeks, preferably between 2 to 8 weeks, more preferably between about 3 to 7 weeks, and even more preferably for about
30 4, 5, or 6 weeks. It will also be appreciated that the effective dosage of a vitamin D₃ compound used for treatment may increase or decrease over the course of a particular treatment.

The term "enantiomers" refers to two stereoisomers of a compound which are non-superimposable mirror images of one another. An equimolar mixture of two enantiomers is called a "racemic mixture" or a "racemate."

The language "genomic" activities or effects of vitamin D₃ is intended to include those activities mediated by the nuclear receptor for 1 α , 25(OH)₂D₃ (VD₃R), e.g., transcriptional activation of target genes.

The term "haloalkyl" is intended to include alkyl groups as defined above that are mono-, di- or polysubstituted by halogen, e.g., fluoromethyl and trifluoromethyl.

The term "halogen" designates -F, -Cl, -Br or -I.

The term "hydroxyl" means -OH.

The term "heteroatom" as used herein means an atom of any element other than carbon or hydrogen. Preferred heteroatoms are nitrogen, oxygen, sulfur and phosphorus.

The term "homeostasis" is art-recognized to mean maintenance of static, or constant, conditions in an internal environment.

The language "hormone secretion" is art-recognized and includes activities of vitamin D₃ compounds that control the transcription and processing responsible for secretion of a given hormone e.g., a parathyroid hormone (PTH) of a vitamin D₃ responsive cell (Bouillon, R. *et al.* (1995) *Endocrine Reviews* 16(2):235-237).

The language "hypercalcemia" or "hypercalcemic activity" is intended to have its accepted clinical meaning, namely, increases in calcium serum levels that are manifested in a subject by the following side effects, depression of central and peripheral nervous system, muscular weakness, constipation, abdominal pain, lack of appetite and, depressed relaxation of the heart during diastole. Symptomatic manifestations of hypercalcemia are triggered by a stimulation of at least one of the following activities, intestinal calcium transport, bone calcium metabolism and osteocalcin synthesis (reviewed in Bouillon, R. *et al.* (1995) *Endocrinology Reviews* 16(2): 200-257).

The terms "hyperproliferative" and "neoplastic" are used interchangeably, and include those cells having the capacity for autonomous growth, i.e., an abnormal state or condition characterized by rapidly proliferating cell growth. Hyperproliferative and neoplastic disease states may be categorized as pathologic, i.e., characterizing or constituting a disease state, or may be categorized as non-pathologic, i.e., a deviation from normal but not associated with a disease state. The term is meant to include all types of cancerous growths or oncogenic processes, metastatic tissues or malignantly

transformed cells, tissues, or organs, irrespective of histopathologic type or stage of invasiveness. "Pathologic hyperproliferative" cells occur in disease states characterized by malignant tumor growth. Examples of non-pathologic hyperproliferative cells include proliferation of cells associated with wound repair.

- 5 The language "immunoglobulin-like transcript 3" or "ILT3" refers to a cell surface molecule of the immunoglobulin superfamily, which is expressed by antigen-presenting cells (APCs) such as monocytes, macrophages and dendritic cells. ILT3 is a member of the immunoglobulin-like transcript (ILT) family and displays a long cytoplasmic tail containing putative immunoreceptor tyrosine-based inhibitory motifs
- 10 (ITIMs). ILT3 has been shown to behave as an inhibitory receptor when cross-linked to a stimulatory receptor. A cytoplasmic component of the ILT3-mediated signaling pathway is the SH2-containing phosphatase SHP-1, which becomes associated with ILT3 upon cross-linking. ILT3 is also internalized and ILT3 ligands are efficiently presented to specific T cells (see, e.g., Cella, M. *et al.* (1997) *J. Exp. Med.* 185:1743).
- 15 The determination of whether the candidate vitamin D₃ compound modulates the expression of the ILT3 surface molecule can be accomplished, for example, by comparison of ILT3 surface molecule expression to a control, by measuring mRNA expression, or by measuring protein expression.

- An "ILT3-associated disorder" includes a disease, disorder or condition which is
- 20 associated with an ILT3 molecule. ILT3 associated disorders include disorders in which ILT3 activity is aberrant or in which a non-ILT3 activity that would benefit from modulation of an ILT3 activity is aberrant. In one embodiment, the ILT3-associated disorder is an immune disorder, e.g., an autoimmune disorder, such as type 1 insulin-dependent diabetes mellitus, adult respiratory distress syndrome, inflammatory bowel
- 25 disease, dermatitis, meningitis, thrombotic thrombocytopenic purpura, Sjogren's syndrome, encephalitis, uveitis, uveoretinitis, leukocyte adhesion deficiency, rheumatoid arthritis, rheumatic fever, Reiter's syndrome, psoriatic arthritis, progressive systemic sclerosis, primary biliary cirrhosis, pemphigus, pemphigoid, necrotizing vasculitis, myasthenia gravis, multiple sclerosis, lupus erythematosus, polymyositis, sarcoidosis,
- 30 granulomatosis, vasculitis, pernicious anemia, CNS inflammatory disorder, antigen-antibody complex mediated diseases, autoimmune haemolytic anemia, Hashimoto's thyroiditis, Graves disease, habitual spontaneous abortions, Reynard's syndrome, glomerulonephritis, dermatomyositis, chronic active hepatitis, celiac disease,

autoimmune complications of AIDS, atrophic gastritis, ankylosing spondylitis and Addison's disease; or transplant rejection, such as GVHD. In certain embodiments of the invention, the ILT3 associated disorder is an immune disorders, such as transplant rejections, graft versus host disease and autoimmune disorders.

5 The term "immune response" includes T and/or B cell responses, e.g., cellular and/or humoral immune responses. The claimed methods can be used to reduce both primary and secondary immune responses. The immune response of a subject can be determined by, for example, assaying antibody production, immune cell proliferation, the release of cytokines, the expression of cell surface markers, cytotoxicity, and the
10 like.

 The terms "immunological tolerance" or "tolerance to an antigen" or "immune tolerance" include unresponsiveness to an antigen without the induction of a prolonged generalized immune deficiency. Consequently, according to the invention, a tolerant host is capable of reacting to antigens other than the tolerizing antigen. Tolerance
15 represents an induced depression in the response of a subject that, had it not been subjected to the tolerance-inducing procedure, would be competent to mount an immune response to that antigen. In one embodiment of the invention, immunological tolerance is induced in an antigen-presenting cell, e.g., an antigen-presenting cell derived from the myeloid or lymphoid lineage, dendritic cells, monocytes and macrophages.

20 The language "immunosuppressive activity" refers to the process of inhibiting a normal immune response. Included in this response is when T and/or B clones of lymphocytes are depleted in size or suppressed in their reactivity, expansion or differentiation. Immunosuppressive activity may be in the form of inhibiting or blocking an immune response already in progress or may involve preventing the
25 induction of an immune response. The functions of activated T cells may be inhibited by suppressing immune cell responses or by inducing specific tolerance, or both. Immunosuppression of T cell responses is generally an active, non-antigen-specific, process that requires continuous exposure of the T cells to the suppressive agent. Tolerance, which involves inducing non-responsiveness or anergy in T cells, is
30 distinguishable from immunosuppression in that it is generally antigen-specific and persists after exposure to the tolerizing agent has ceased. Operationally, tolerance can be demonstrated by the lack of a T cell response upon re-exposure to specific antigen in the absence of the tolerizing agent.

The language "improved biological properties" refers to any activity inherent in a compound of the invention that enhances its effectiveness *in vivo*. In a preferred embodiment, this term refers to any qualitative or quantitative improved therapeutic property of a vitamin D₃ compound, such as reduced toxicity, *e.g.*, reduced

5 hypercalcemic activity.

The language "inhibiting the growth" of the neoplasm includes the slowing, interrupting, arresting or stopping its growth and metastases and does not necessarily indicate a total elimination of the neoplastic growth.

The phrase "inhibition of an immune response" is intended to include decreases
10 in T cell proliferation and activity, *e.g.*, a decrease in IL₂, interferon- γ , GM-CSF synthesis and secretion (Lemire, J. M. (1992) *J. Cell Biochemistry* 49:26-31, Lemire, J. M. *et al.* (1994) *Endocrinology* 135 (6): 2813-2821; Bouillon, R. *et al.* (1995) *Endocrine Review* 16 (2):231-32).

The term "isomers" or "stereoisomers" refers to compounds which have identical
15 chemical constitution, but differ with regard to the arrangement of the atoms or groups in space.

The term "leukemia" is intended to have its clinical meaning, namely, a neoplastic disease in which white corpuscle maturation is arrested at a primitive stage of cell development. The disease is characterized by an increased number of leukemic
20 blast cells in the bone marrow, and by varying degrees of failure to produce normal hematopoietic cells. The condition may be either acute or chronic. Leukemias are further typically categorized as being either lymphocytic *i.e.*, being characterized by cells which have properties in common with normal lymphocytes, or myelocytic (or myelogenous), *i.e.*, characterized by cells having some characteristics of normal
25 granulocytic cells. Acute lymphocytic leukemia ("ALL") arises in lymphoid tissue, and ordinarily first manifests its presence in bone marrow. Acute myelocytic leukemia ("AML") arises from bone marrow hematopoietic stem cells or their progeny. The term acute myelocytic leukemia subsumes several subtypes of leukemia: myeloblastic leukemia, promyelocytic leukemia, and myelomonocytic leukemia. In addition,
30 leukemias with erythroid or megakaryocytic properties are considered myelogenous leukemias as well.

The term "leukemic cancer" refers to all cancers or neoplasias of the hemopoietic and immune systems (blood and lymphatic system). The acute and chronic leukemias,

together with the other types of tumors of the blood, bone marrow cells (myelomas), and lymph tissue (lymphomas), cause about 10% of all cancer deaths and about 50% of all cancer deaths in children and adults less than 30 years old. Chronic myelogenous leukemia (CML), also known as chronic granulocytic leukemia (CGL), is a neoplastic disorder of the hematopoietic stem cell. The term "leukemia" is art recognized and refers to a progressive, malignant disease of the blood-forming organs, marked by distorted proliferation and development of leukocytes and their precursors in the blood and bone marrow.

The term "modulate" refers to increases or decreases in the activity of a cell in response to exposure to a compound of the invention, *e.g.*, the inhibition of proliferation and/or induction of differentiation of at least a sub-population of cells in an animal such that a desired end result is achieved, *e.g.*, a therapeutic result. In preferred embodiments, this phrase is intended to include hyperactive conditions that result in pathological disorders.

The common medical meaning of the term "neoplasia" refers to "new cell growth" that results as a loss of responsiveness to normal growth controls, *e.g.* to neoplastic cell growth. A "hyperplasia" refers to cells undergoing an abnormally high rate of growth. However, as used herein, the terms neoplasia and hyperplasia can be used interchangeably, as their context will reveal, referring to generally to cells experiencing abnormal cell growth rates. Neoplasias and hyperplasias include "tumors," which may be either benign, premalignant or malignant.

The language "non-genomic" vitamin D₃ activities include cellular (*e.g.*, calcium transport across a tissue) and subcellular activities (*e.g.*, membrane calcium transport opening of voltage-gated calcium channels, changes in intracellular second messengers) elicited by vitamin D₃ compounds in a responsive cell. Electrophysiological and biochemical techniques for detecting these activities are known in the art. An example of a particular well-studied non-genomic activity is the rapid hormonal stimulation of intestinal calcium mobilization, termed "transcaltachia" (Nemere I. *et al.* (1984) *Endocrinology* 115:1476-1483; Lieberherr M. *et al.* (1989) *J. Biol. Chem.* 264:20403-20406; Wali R.K. *et al.* (1992) *Endocrinology* 131:1125-1133; Wali R.K. *et al.* (1992) *Am. J. Physiol.* 262:G945-G953; Wali R.K. *et al.* (1990) *J. Clin. Invest.* 85:1296-1303; Bolt M.J.G. *et al.* (1993) *Biochem. J.* 292:271-276). Detailed descriptions of experimental transcaltachia are provided in Norman, A.W. (1993) *Endocrinology*

268(27):20022-20030; Yoshimoto, Y. and Norman, A.W. (1986)

Endocrinology 118:2300-2304. Changes in calcium activity and second messenger systems are well known in the art and are extensively reviewed in Bouillion, R. *et al.* (1995) *Endocrinology Review* 16(2): 200-257; the description of which is incorporated
5 herein by reference.

The phrases "parenteral administration" and "administered parenterally" as used herein means modes of administration other than enteral and topical administration, usually by injection, and includes, without limitation, intravenous, intramuscular, intraarterial, intrathecal, intracapsular, intraorbital, intracardiac, intradermal,
10 intraperitoneal, transtracheal, subcutaneous, subcuticular, intraarticular, subcapsular, subarachnoid, intraspinal and intrasternal injection and infusion.

The terms "polycyclyl" or "polycyclic radical" refer to the radical of two or more cyclic rings (*e.g.*, cycloalkyls, cycloalkenyls, cycloalkynyls, aryls and/or heterocyclyls) in which two or more carbons are common to two adjoining rings, *e.g.*, the rings are
15 "fused rings". Rings that are joined through non-adjacent atoms are termed "bridged" rings. Each of the rings of the polycycle can be substituted with such substituents as described above, as for example, halogen, hydroxyl, alkylcarbonyloxy, arylcarbonyloxy, alkoxycarbonyloxy, aryloxy carbonyloxy, carboxylate, alkylcarbonyl, alkoxycarbonyl, aminocarbonyl, alkylthiocarbonyl, alkoxyl, phosphate, phosphonato, phosphinato,
20 cyano, amino (including alkyl amino, dialkylamino, arylamino, diarylamino, and alkylaryl amino), acylamino (including alkylcarbonylamino, arylcarbonylamino, carbamoyl and ureido), amidino, imino, sulfhydryl, alkylthio, arylthio, thiocarboxylate, sulfates, sulfonato, sulfamoyl, sulfonamido, nitro, trifluoromethyl, cyano, azido, heterocyclyl, alkyl, alkylaryl, or an aromatic or heteroaromatic moiety.

25 The term "prodrug" includes compounds with moieties which can be metabolized *in vivo*. Generally, the prodrugs are metabolized *in vivo* by esterases or by other mechanisms to active drugs. Examples of prodrugs and their uses are well known in the art (See, *e.g.*, Berge *et al.* (1977) "Pharmaceutical Salts", *J. Pharm. Sci.* 66:1-19). The prodrugs can be prepared *in situ* during the final isolation and purification of the
30 compounds, or by separately reacting the purified compound in its free acid form or hydroxyl with a suitable esterifying agent. Hydroxyl groups can be converted into esters *via* treatment with a carboxylic acid. Examples of prodrug moieties include substituted and unsubstituted, branch or unbranched lower alkyl ester moieties, (*e.g.*, propionol

acid esters), lower alkenyl esters, di-lower alkyl-amino lower-alkyl esters (*e.g.*, dimethylaminoethyl ester), acylamino lower alkyl esters (*e.g.*, acetyloxymethyl ester), acyloxy lower alkyl esters (*e.g.*, pivaloyloxymethyl ester), aryl esters (phenyl ester), aryl-lower alkyl esters (*e.g.*, benzyl ester), substituted (*e.g.*, with methyl, halo, or methoxy substituents) aryl and aryl-lower alkyl esters, amides, lower-alkyl amides, di-lower alkyl amides, and hydroxy amides. Preferred prodrug moieties are propionoic acid esters and acyl esters. Prodrugs which are converted to active forms through other mechanisms *in vivo* are also included.


The language "a prophylactically effective anti-neoplastic amount" of a compound refers to an amount of a vitamin D₃ compound of the formula (I) or otherwise described herein which is effective, upon single or multiple dose administration to the patient, in preventing or delaying the occurrence of the onset of a neoplastic disease state.

The term "psoriasis" is intended to have its medical meaning, namely, a disease which afflicts primarily the skin and produces raised, thickened, scaling, nonscarring lesions. The lesions are usually sharply demarcated erythematous papules covered with overlapping shiny scales. The scales are typically silvery or slightly opalescent. Involvement of the nails frequently occurs resulting in pitting, separation of the nail, thickening and discoloration. Psoriasis is sometimes associated with arthritis, and it may be crippling.

The language "reduced toxicity" is intended to include a reduction in any undesired side effect elicited by a vitamin D₃ compound when administered *in vivo*, *e.g.*, a reduction in the hypercalcemic activity.

The term "sarcoma" is art recognized and refers to malignant tumors of mesenchymal derivation.

The term "secosteroid" is art-recognized and includes compounds in which one of the cyclopentanoperhydro-phenanthrene rings of the steroid ring structure is broken. 1 α ,25(OH)₂D₃ and analogs thereof are hormonally active secosteroids. In the case of vitamin D₃, the 9-10 carbon-carbon bond of the B-ring is broken, generating a seco-B-steroid. The official IUPAC name for vitamin D₃ is 9,10-secocholesta-5,7,10(19)-trien-3B-ol. For convenience, a 6-*s-trans* conformer of 1 α ,25(OH)₂D₃ is illustrated herein having all carbon atoms numbered using standard steroid notation.



In the formulas presented herein, the various substituents on ring A are illustrated as joined to the steroid nucleus by one of these notations: a dotted line (—) indicating a substituent which is in the β -orientation (*i.e.*, above the plane of the ring), a wedged solid line (\blacktriangleleft) indicating a substituent which is in the α -orientation (*i.e.*, below the plane of the molecule), or a wavy line (\sim) indicating that a substituent may be either above or below the plane of the ring. In regard to ring A, it should be understood that the stereochemical convention in the vitamin D field is opposite from the general chemical field, wherein a dotted line indicates a substituent on Ring A which is in an α -orientation (*i.e.*, below the plane of the molecule), and a wedged solid line indicates a substituent on ring A which is in the β -orientation (*i.e.*, above the plane of the ring). As shown, the A ring of the hormone $1\alpha,25(\text{OH})_2\text{D}_3$ contains two asymmetric centers at carbons 1 and 3, each one containing a hydroxyl group in well-characterized configurations, namely the 1α - and 3β - hydroxyl groups. In other words, carbons 1 and 3 of the A ring are said to be "chiral carbons" or "carbon centers."

The term "sulfhydryl" or "thiol" means $-\text{SH}$.

The term "subject" includes organisms which are capable of suffering from a vitamin D_3 associated state or who could otherwise benefit from the administration of a vitamin D_3 compound of the invention, such as human and non-human animals. Preferred human animals include human patients suffering from or prone to suffering from a vitamin D_3 associated state, as described herein. The term "non-human animals" of the invention includes all vertebrates, *e.g.*, mammals, *e.g.*, rodents, *e.g.*, mice, and

non-mammals, such as non-human primates, sheep, dog, cow, chickens, amphibians, reptiles, etc.

The phrases "systemic administration," "administered systemically", "peripheral administration" and "administered peripherally" as used herein mean the administration of a vitamin D₃ compound(s), drug or other material, such that it enters the patient's system and, thus, is subject to metabolism and other like processes, for example, subcutaneous administration.

The language "therapeutically effective anti-neoplastic amount" of a vitamin D₃ compound of the invention refers to an amount of an agent which is effective, upon single or multiple dose administration to the patient, in inhibiting the growth of a neoplastic vitamin D₃-responsive cells, or in prolonging the survivability of the patient with such neoplastic cells beyond that expected in the absence of such treatment.

The language "transplant rejection" refers to an immune reaction directed against a transplanted organ(s) from other human donors (allografts) or from other species such as sheep, pigs, or non-human primates (xenografts). Therefore, the method of the invention is useful for preventing an immune reaction to transplanted organs from other human donors (allografts) or from other species (xenografts). Such tissues for transplantation include, but are not limited to, heart, liver, kidney, lung, pancreas, pancreatic islets, bone marrow, brain tissue, cornea, bone, intestine, skin, and hematopoietic cells. Also included within this definition is "graft versus host disease" of "GVHD," which is a condition where the graft cells mount an immune response against the host. Therefore, the method of the invention is useful in preventing graft versus host disease in cases of mismatched bone marrow or lymphoid tissue transplanted for the treatment of acute leukemia, aplastic anemia, and enzyme or immune deficiencies, for example. The term "transplant rejection" also includes disease symptoms characterized by loss of organ function. For example, kidney rejection would be characterized by a rising creatine level in blood. Heart rejection is characterized by an endomyocardial biopsy, while pancreas rejection is characterized by rising blood glucose levels. Liver rejection is characterized by the levels of transaminases of liver origin and bilirubin levels in blood. Intestine rejection is determined by biopsy, while lung rejection is determined by measurement of blood oxygenation.

The terms "urogenital", "urogenital system" and "urogenital tract" are used interchangeably and are intended to include all organs involved in reproduction and in

the formation and voidance of urine. Included with in these terms are the kidneys, bladder and prostate. The term "VDR" is intended to include members of the type II class of steroid/thyroid superfamily of receptors (Stunnenberg, H.G. (1993) *Bio Essays* 15(5):309-15), which are able to bind and transactivate through the vitamin D response element (VDRE) in the absence of a ligand (Damm *et al.* (1989) *Nature* 339:593-97; Sap *et al.* *Nature* 343:177-180).

The term "VDRE" refers to DNA sequences composed of half-sites arranged as direct repeats. It is known in the art that type II receptors do not bind to their respective binding site as homodimers but require an auxiliary factor, RXR (*e.g.* RXR α , RXR β , RXR γ) for high affinity binding Yu *et al.* (1991) *Cell* 67:1251-1266; Bugge *et al.* (1992) *EMBO J.* 11:1409-1418; Kliewer *et al.* (1992) *Nature* 355:446-449; Leid *et al.* (1992) *EMBO J.* 11:1419-1435; Zhang *et al.* (1992) *Nature* 355:441-446).

The language "vitamin D₃ associated state" is a state which can be prevented, treated or otherwise ameliorated by administration of one or more compounds of the invention. Vitamin D₃ associated states include ILT3-associated disorders, disorders characterized by an aberrant activity of a vitamin D₃-responsive cell, disorders characterized by a deregulation of calcium and phosphate metabolism, and other disorders or states described herein.


The term "vitamin D₃-responsive cell" includes any cell which is capable of responding to a vitamin D₃ compound having the formula I or otherwise described herein, or is associated with disorders involving an aberrant activity of hyperproliferative skin cells, parathyroid cells, neoplastic cells, immune cells, and bone cells. These cells can respond to vitamin D₃ activation by triggering genomic and/or non-genomic responses that ultimately result in the modulation of cell proliferation, differentiation survival, and/or other cellular activities such as hormone secretion. In a preferred embodiment, the ultimate responses of a cell are inhibition of cell proliferation and/or induction of differentiation-specific genes. Exemplary vitamin D₃ responsive cells include immune cells, bone cells, neuronal cells, endocrine cells, neoplastic cells, epidermal cells, endodermal cells, smooth muscle cells, among others.

With respect to the nomenclature of a chiral center, terms "d" and "l" configuration are as defined by the IUPAC Recommendations. As to the use of the terms, diastereomer, racemate, epimer and enantiomer will be used in their normal context to describe the stereochemistry of preparations.


2. VITAMIN D₃ COMPOUNDS OF THE INVENTION

- 5 The acylated vitamin D₃ compounds of formula I above exert a full spectrum of 1,25(OH)₂D₃ biological activities such as binding to the specific nuclear receptor VDR, suppression of the increased parathyroid hormone levels in 5,6-nephrectomized rats, suppression of INF-γ release in MLR cells, stimulation of HL-60 leukemia cell differentiation and inhibition of solid tumor cell proliferation. It is well known that *in*
- 10 *vivo* and in cellular cultures 1,25-(OH)₂D₃ undergoes a cascade of metabolic modifications initiated by the influence of 24R-hydroxylase enzyme. First 24R-hydroxy metabolite is formed, which is oxydized to 24-keto intermediate, and then 23S-hydroxylation and fragmentation produce the fully inactive calcitroic acid.

- Unexpectedly, it has been found that 1,3-O-diacetyl-1,25-dihydroxy-16-ene-24-
- 15 keto-19-nor-cholecalciferol, a compound in accordance with the invention (shown below), has a maximum tolerated dose (MTD) that is more than 300 times that of the corresponding dihydroxy compound, 1,25-dihydroxy-16-ene-24keto-19-nor-cholecalciferol (also shown below).




20 1,3-O-Diacetyl-1,25-dihydroxy-16-ene-24-keto-19-nor-cholecalciferol



1,25-dihydroxy-16-ene-24-keto-19-nor-cholecalciferol

Thus, in one aspect, the invention provides vitamin D₃ compounds of the formula I:



wherein:

X_1 and X_2 are each independently H_2 or $=CH_2$, provided X_1 and X_2 are not both $=CH_2$;

R_1 and R_2 are each independently, hydroxyl, $OC(O)C_1-C_4$ alkyl, $OC(O)$ hydroxyalkyl, $OC(O)$ fluoroalkyl, provided that R_1 and R_2 are not both hydroxyl;


R_3 and R_4 are each independently hydrogen, C_1-C_4 alkyl, hydroxyalkyl, or haloalkyl, or R_3 and R_4 taken together with C_{20} form C_3-C_6 cycloalkyl; and

R_5 and R_6 are each independently C_1-C_4 alkyl, hydroxyalkyl, or haloalkyl; and pharmaceutically acceptable esters, salts, and prodrugs thereof.

In one embodiment, X_1 and X_2 are each H_2 . In another embodiment, R_3 is hydrogen and R_4 is C_1-C_4 alkyl. In a preferred embodiment R_4 is methyl.

In another embodiment, R_5 and R_6 are each independently methyl, ethyl or fluoromethyl. In a preferred embodiment, R_5 and R_6 are each methyl.

In yet another embodiment, R_1 and R_2 are each independently hydroxyl or $OC(O)C_1-C_4$ alkyl. In a preferred embodiment, R_1 and R_2 are each $OC(O)C_1-C_4$ alkyl. In another preferred embodiment, R_1 and R_2 are each acetyloxy. In still another embodiment, the haloalkyl group can be a fluoroalkyl group, for example, fluoromethyl and trifluoromethyl. A preferred compound is 1,3-O-diacetyl-1,25-dihydroxy-16-ene-24-keto-19-nor-cholecalciferol, having the following structure:



5

The structures of some of the compounds of the invention include asymmetric
10 carbon atoms. Accordingly, the isomers arising from such asymmetry (*e.g.*, all
enantiomers and diastereomers) are included within the scope of the invention, unless
indicated otherwise. Such isomers can be obtained in substantially pure form by
classical separation techniques and/or by stereochemically controlled synthesis.

Naturally occurring or synthetic isomers can be separated in several ways known
15 in the art. Methods for separating a racemic mixture of two enantiomers include
chromatography using a chiral stationary phase (see, *e.g.*, "Chiral Liquid
Chromatography," W.J. Lough, Ed. Chapman and Hall, New York (1989)).
Enantiomers can also be separated by classical resolution techniques. For example,
formation of diastereomeric salts and fractional crystallization can be used to separate
20 enantiomers. For the separation of enantiomers of carboxylic acids, the diastereomeric
salts can be formed by addition of enantiomerically pure chiral bases such as brucine,
quinine, ephedrine, strychnine, and the like. Alternatively, diastereomeric esters can be
formed with enantiomerically pure chiral alcohols such as menthol, followed by
separation of the diastereomeric esters and hydrolysis to yield the free, enantiomerically

enriched carboxylic acid. For separation of the optical isomers of amino compounds, addition of chiral carboxylic or sulfonic acids, such as camphorsulfonic acid, tartaric acid, mandelic acid, or lactic acid can result in formation of the diastereomeric salts.

5 3. USES OF THE VITAMIN D₃ COMPOUNDS OF THE INVENTION

 In another embodiment, the invention also provides methods for treating a subject for a vitamin D₃ associated state, by administering to the subject an effective amount of a vitamin D₃ compound of formula I or otherwise described herein. Vitamin D₃ associated states include disorders involving an aberrant activity of a vitamin D₃-responsive cell, e.g., neoplastic cells, hyperproliferative skin cells, parathyroid cells, 10 immune cells and bone cells, among others. Vitamin D₃ associated states also include ILT3-associated disorders. In certain embodiments, the subject is a mammal, e.g., a primate, e.g., a human.

 In certain embodiments, the methods of the invention include administering to a 15 subject a therapeutically effective amount of a vitamin D₃ compound in combination with another pharmaceutically active compound. Examples of pharmaceutically active compounds include compounds known to treat autoimmune disorders, e.g., immunosuppressant agents such as cyclosporin A, rapamycin, desoxyspergualine, FK 506, steroids, azathioprine, anti-T cell antibodies and monoclonal antibodies to T cell 20 subpopulations. Other pharmaceutically active compounds that may be used can be found in *Harrison's Principles of Internal Medicine*, Thirteenth Edition, Eds. T.R. Harrison *et al.* McGraw-Hill N.Y., NY; and the Physicians Desk Reference 50th Edition 1997, Oradell New Jersey, Medical Economics Co., the complete contents of which are expressly incorporated herein by reference. The angiogenesis inhibitor compound and 25 the pharmaceutically active compound may be administered to the subject in the same pharmaceutical composition or in different pharmaceutical compositions (at the same time or at different times).

A. Hyperproliferative Conditions

30 In another aspect, the present invention provides a method of treating a subject for a disorder characterized by aberrant activity of a vitamin D₃-responsive cell. The method involves administering to the subject an effective amount of a pharmaceutical

composition of a vitamin D₃ compound of formula I or otherwise described herein such that the activity of the cell is modulated.

In certain embodiments, the cells to be treated are hyperproliferative cells. As described in greater detail below, the vitamin D₃ compounds of the invention can be used to inhibit the proliferation of a variety of hyperplastic and neoplastic tissues. In accordance with the present invention, vitamin D₃ compounds of the invention can be used in the treatment of both pathologic and non-pathologic proliferative conditions characterized by unwanted growth of vitamin D₃-responsive cells, *e.g.*, hyperproliferative skin cells, immune cells, and tissue having transformed cells, *e.g.*, such as carcinomas, sarcomas and leukemias. In other embodiments, the cells to be treated are aberrant secretory cells, *e.g.*, parathyroid cells, immune cells.

The use of vitamin D compounds in treating hyperproliferative conditions has been limited because of their hypercalcemic effects. Thus, vitamin D₃ compounds of the invention can provide a less toxic alternative to current methods of treatment.

In one embodiment, the invention features a method for inhibiting the proliferation and/or inducing the differentiation of a hyperproliferative skin cell, *e.g.*, an epidermal or an epithelial cell, *e.g.*, a keratinocytes, by contacting the cells with a vitamin D₃ compound of the invention. In general, the method includes a step of contacting a pathological or non-pathological hyperproliferative cell with an effective amount of such vitamin D₃ compound to promote the differentiation of the hyperproliferative cells. The present method can be performed on cells in culture, *e.g.*, *in vitro* or *ex vivo*, or can be performed on cells present in an animal subject, *e.g.*, as part of an *in vivo* therapeutic protocol. The therapeutic regimen can be carried out on a human or any other animal subject.

The vitamin D₃ compounds of the present invention can be used to treat a hyperproliferative skin disorder. Exemplary disorders include, but are not limited to, psoriasis, basal cell carcinoma, keratinization disorders and keratosis. Additional examples of these disorders include eczema; lupus associated skin lesions; psoriatic arthritis; rheumatoid arthritis that involves hyperproliferation and inflammation of epithelial-related cells lining the joint capsule; dermatitides such as seborrheic dermatitis and solar dermatitis; keratoses such as seborrheic keratosis, senile keratosis, actinic keratosis, photo-induced keratosis, and keratosis follicularis; acne vulgaris; keloids and prophylaxis against keloid formation; nevi; warts including verruca, condyloma or

condyloma acuminatum, and human papilloma viral (HPV) infections such as venereal warts; leukoplakia; lichen planus; and keratitis.

- In an illustrative example, vitamin D₃ compounds of the invention can be used to inhibit the hyperproliferation of keratinocytes in treating diseases such as psoriasis by administering an effective amount of these compounds to a subject in need of treatment. The term "psoriasis" is intended to have its medical meaning, namely, a disease which afflicts primarily the skin and produces raised, thickened, scaling, nonscarring lesions. The lesions are usually sharply demarcated erythematous papules covered with overlapping shiny scales. The scales are typically silvery or slightly opalescent.
- 10 Involvement of the nails frequently occurs resulting in pitting, separation of the nail, thickening and discoloration. Psoriasis is sometimes associated with arthritis, and it may be crippling. Hyperproliferation of keratinocytes is a key feature of psoriatic epidermal hyperplasia along with epidermal inflammation and reduced differentiation of keratinocytes. Multiple mechanisms have been invoked to explain the keratinocyte
- 15 hyperproliferation that characterizes psoriasis. Disordered cellular immunity has also been implicated in the pathogenesis of psoriasis.

B. Neoplasia

- The invention also features methods for inhibiting the proliferation and/or reversing the transformed phenotype of vitamin D₃-responsive hyperproliferative cells by contacting the cells with a vitamin D₃ compound of formula I or otherwise described herein. In general, the method includes a step of contacting pathological or non-pathological hyperproliferative cells with an effective amount of a vitamin D₃ compound of the invention for promoting the differentiation of the hyperproliferative cells. The
- 20 present method can be performed on cells in culture, *e.g.*, *in vitro* or *ex vivo*, or can be performed on cells present in an animal subject, *e.g.*, as part of an *in vivo* therapeutic protocol. The therapeutic regimen can be carried out on a human or other subject.

- The vitamin D₃ compounds of formula I or otherwise described herein can be tested initially *in vitro* for their inhibitory effects in the proliferation of neoplastic cells.
- 30 Examples of cell lines that can be used are transformed cells, *e.g.*, the human promyeloid leukemia cell line HL-60, and the human myeloid leukemia U-937 cell line (Abe E. *et al.* (1981) *Proc. Natl. Acad. Sci. USA* 78:4990-4994; Song L.N. and Cheng T. (1992) *Biochem Pharmacol* 43:2292-2295; Zhou J.Y. *et al.* (1989) *Blood* 74:82-93; U.S.

Pat. Nos. 5,401,733, U.S. 5,087,619). Alternatively, the antitumoral effects of vitamin D₃ compounds of the invention can be tested *in vivo* using various animal models known in the art and summarized in Bouillon, R. *et al.* (1995) *Endocrine Reviews* 16(2):233 (Table E), which is incorporated by reference herein. For example, SL mice are
5 routinely used in the art to test vitamin D compounds as models for M1 myeloid leukemia (Honma *et al.* (1983) *Cell Biol.* 80:201-204; Kasukabe T. *et al.* (1987) *Cancer Res.* 47:567-572); breast cancer studies can be performed in, for example, nude mice models for human MX1 (ER) (Abe J. *et al.* (1991) *Endocrinology* 129:832-837; other cancers, *e.g.*, colon cancer, melanoma osteosarcoma, can be characterized in, for
10 example, nude mice models as describe in (Eisman J. A. *et al.* (1987) *Cancer Res.* 47:21-25; Kawaura A. *et al.* (1990) *Cancer Lett* 55:149-152; Belleli A. (1992) *Carcinogenesis* 13:2293-2298; Tsuchiya H. *et al.* (1993) *J. Orthopaed Res.* 11:122-130).

The subject method may also be used to inhibit the proliferation of hyperplastic/neoplastic cells of hematopoietic origin, *e.g.*, arising from myeloid,
15 lymphoid or erythroid lineages, or precursor cells thereof. For instance, the present invention contemplates the treatment of various myeloid disorders including, but not limited to, acute promyeloid leukemia (APML), acute myelogenous leukemia (AML) and chronic myelogenous leukemia (CML) (reviewed in Vaickus, L. (1991) *Crit Rev. in Oncol/Hemotol.* 11:267-97). Lymphoid malignancies which may be treated by the
20 subject method include, but are not limited to acute lymphoblastic leukemia (ALL) which includes B-lineage ALL and T-lineage ALL, chronic lymphocytic leukemia (CLL), prolymphocytic leukemia (PLL), hairy cell leukemia (HLL) and Waldenstrom's macroglobulinemia (WM). Additional forms of malignant lymphomas contemplated by the treatment method of the present invention include, but are not limited to non-
25 Hodgkin lymphoma and variants thereof, peripheral T cell lymphomas, adult T cell leukemia/lymphoma (ATL), cutaneous T-cell lymphoma (CTCL), large granular lymphocytic leukemia (LGF) and Hodgkin's disease.

In certain embodiments, the vitamin D₃ compounds of the invention can be used in combinatorial therapy with conventional cancer chemotherapeutics. Conventional
30 treatment regimens for leukemia and for other tumors include radiation, drugs, or a combination of both. In addition to radiation, the following drugs, usually in combinations with each other, are often used to treat acute leukemias: vincristine, prednisone, methotrexate, mercaptopurine, cyclophosphamide, and cytarabine. In

chronic leukemia, for example, busulfan, melphalan, and chlorambucil can be used in combination. All of the conventional anti-cancer drugs are highly toxic and tend to make patients quite ill while undergoing treatment. Vigorous therapy is based on the premise that unless every leukemic cell is destroyed, the residual cells will multiply and
5 cause a relapse.

The subject method can also be useful in treating malignancies of the various organ systems, such as affecting lung, breast, lymphoid, gastrointestinal, and genito-urinary tract as well as adenocarcinomas which include malignancies such as most colon cancers, renal-cell carcinoma, prostate cancer and/or testicular tumors, non-small cell
10 carcinoma of the lung, cancer of the small intestine and cancer of the esophagus.

According to the general paradigm of vitamin D₃ involvement in differentiation of transformed cells, exemplary solid tumors that can be treated according to the method of the present invention include vitamin D₃-responsive phenotypes of sarcomas and carcinomas such as, but not limited to: fibrosarcoma, myxosarcoma, liposarcoma,
15 chondrosarcoma, osteogenic sarcoma, chordoma, angiosarcoma, endotheliosarcoma, lymphangiosarcoma, lymphangioendotheliosarcoma, synovioma, mesothelioma, Ewing's tumor, leiomyosarcoma, rhabdomyosarcoma, colon carcinoma, pancreatic cancer, breast cancer, ovarian cancer, prostate cancer, squamous cell carcinoma, basal cell carcinoma, adenocarcinoma, sweat gland carcinoma, sebaceous gland carcinoma,
20 papillary carcinoma, papillary adenocarcinomas, cystadenocarcinoma, medullary carcinoma, bronchogenic carcinoma, renal cell carcinoma, hepatoma, bile duct carcinoma, choriocarcinoma, seminoma, embryonal carcinoma, Wilms' tumor, cervical cancer, testicular tumor, lung carcinoma, small cell lung carcinoma, bladder carcinoma, epithelial carcinoma, glioma, astrocytoma, medulloblastoma, craniopharyngioma,
25 ependymoma, pinealoma, hemangioblastoma, acoustic neuroma, oligodendroglioma, meningioma, melanoma, neuroblastoma, and retinoblastoma.

Determination of a therapeutically effective anti-neoplastic amount or a prophylactically effective anti-neoplastic amount of the vitamin D₃ compound of the invention, can be readily made by the physician or veterinarian (the "attending
30 clinician"), as one skilled in the art, by the use of known techniques and by observing results obtained under analogous circumstances. The dosages may be varied depending upon the requirements of the patient in the judgment of the attending clinician, the severity of the condition being treated and the particular compound being employed. In

determining the therapeutically effective antineoplastic amount or dose, and the prophylactically effective antineoplastic amount or dose, a number of factors are considered by the attending clinician, including, but not limited to: the specific hyperplastic/neoplastic cell involved; pharmacodynamic characteristics of the particular agent and its mode and route of administration; the disorder time course of treatment; the species of mammal; its size, age, and general health; the specific disease involved; the degree of involvement or the severity of the disease; the response of the individual patient; the particular compound administered; the mode of administration; the bioavailability characteristics of the preparation administered; the dose regimen selected; the kind of concurrent treatment (*i.e.*, the interaction of the vitamin D₃ compounds of the invention with other co-administered therapeutics); and other relevant circumstances. U.S. Patent 5,427,916, for example, describes method for predicting the effectiveness of antineoplastic therapy in individual patients, and illustrates certain methods which can be used in conjunction with the treatment protocols of the instant invention.

Treatment can be initiated with smaller dosages which are less than the optimum dose of the compound. Thereafter, the dosage should be increased by small increments until the optimum effect under the circumstances is reached. For convenience, the total daily dosage may be divided and administered in portions during the day if desired. A therapeutically effective antineoplastic amount and a prophylactically effective antineoplastic amount of a vitamin D₃ compound of the invention is expected to vary from about 0.1 milligram per kilogram of body weight per day (mg/kg/day) to about 100 mg/kg/day.

Compounds which are determined to be effective for the prevention or treatment of tumors in animals, *e.g.*, dogs, rodents, may also be useful in treatment of tumors in humans. Those skilled in the art of treating tumors in humans will know, based upon the data obtained in animal studies, the dosage and route of administration of the compound to humans. In general, the dosage and route of administration in humans is expected to be similar to that in animals.

The identification of those patients who are in need of prophylactic treatment for hyperplastic/neoplastic disease states is well within the ability and knowledge of one skilled in the art. Certain of the methods for identification of patients which are at risk of developing neoplastic disease states which can be treated by the subject method are

appreciated in the medical arts, such as family history of the development of a particular disease state and the presence of risk factors associated with the development of that disease state in the subject patient. A clinician skilled in the art can readily identify such candidate patients, by the use of, for example, clinical tests, physical examination and
5 medical/family history.

C. Immunological Activity

Healthy individuals protect themselves against foreign invaders using many different mechanisms, including physical barriers, phagocytic cells in the blood and
10 tissues, a class of immune cells known as lymphocytes, and various blood-born molecules. All of these mechanisms participate in defending individuals from a potentially hostile environment. Some of these defense mechanisms, known as natural or innate immunity, are present in an individual prior to exposure to infectious microbes or other foreign macromolecules, are not enhanced by such exposures, and do not
15 discriminate among most foreign substances. Other defense mechanisms, known as acquired or specific immunity, are induced or stimulated by exposure of foreign substances, are exquisitely specific for distinct macromolecules, and increase in magnitude and defensive capabilities with each successive exposure to a particular macromolecule. Substances that induce a specific immune response are known as
20 antigens (see, e.g., Abbas, A. *et al.*, *Cellular and Molecular Immunology*, W.B. Saunders Company, Philadelphia, 1991; Silverstein, A.M. *A history of Immunology*, San Diego, Academic Press, 1989; Unanue A. *et al.*, *Textbook of Immunology*, 2nd ed. Williams and Wilkens, Baltimore, 1984).

One of the most remarkable properties of the immune system is its ability to
25 distinguish between foreign antigens and self-antigens. Therefore, the lymphocytes in each individual are able to recognize and respond to many foreign antigens but are normally unresponsive to the potentially antigenic substances present in the individual. This immunological unresponsiveness is referred to as immune tolerance (see, e.g., Burt RK *et al.* (2002) *Blood* 99:768; Coutinho, A. *et al.* (2001) *Immunol. Rev.* 182:89;
30 Schwartz, RH (1990) *Science* 248:1349; Miller, J.F. *et al.* (1989) *Immunology Today* 10:53).

Self-tolerance is an acquired process that has to be learned by the lymphocytes of each individual. It occurs in part because lymphocytes pass through a stage in their

development when an encounter with antigen presented by antigen-presenting cells (APCs) leads to their death or inactivation in a process known as positive and negative selection (see, e.g., Debatin KM (2001) *Ann. Hematol.* 80 Suppl 3:B29; Abbas, A. (1991), *supra*). Thus, potentially self-recognizing lymphocytes come into contact with self-antigens at this stage of functional immaturity and are prevented from developing to a stage at which they would be able to respond to self-antigens. Autoimmunity arises when abnormalities in the induction or maintenance of self-tolerance occur that result in a loss of tolerance to a particular antigen(s) and a subsequent attack by the host's immune system on the host's tissues that express the antigen(s) (see, e.g., Boyton RJ *et al.* (2002) *Clin. Exp. Immunol.* 127:4; Hagiwara E. (2001) *Ryumachi* 41:888; Burt RK *et al.* (2002) *Blood* 99:768).

The ability of the immune system to distinguish between self and foreign antigens also plays a critical role in tissue transplantation. The success of a transplant depends on preventing the immune system of the host recipient from recognizing the transplant as foreign and, in some cases, preventing the graft from recognizing the host tissues as foreign. For example, when a host receives a bone marrow transplant, the transplanted bone marrow may recognize the new host as foreign, resulting in graft versus host disease (GVHD). Consequently, the survival of the host depends on preventing both the rejection of the donor marrow as well as rejection of the host by the graft immune reaction (see, e.g., Waldmann H *et al.* (2001) *Int. Arch. Allergy Immunol.* 126:11).

Currently, deleterious immune reactions that result in autoimmune diseases and transplant rejections are prevented or treated using agents such as steroids, azathioprine, anti-T cell antibodies, and more recently, monoclonal antibodies to T cell subpopulations. Immunosuppressive drugs such as cyclosporin A (CsA), rapamycin, desoxyspergualine and FK-506 are also widely used.

Nonspecific immune suppression agents, such as steroids and antibodies to lymphocytes, put the host at increased risk for opportunistic infection and development of tumors. Moreover, many immunosuppressive drugs result in bone demineralization within the host (see, e.g., Chhajed PN *et al.* (2002) *Indian J. Chest Dis. Allied* 44:31; Wijdicks EF (2001) *Liver Transpl.* 7:937; Karamchic J *et al.* (2001) *Med. Arch.* 55:243; U.S. Patent No. 5,597,563 issued to Beschoner, WE and U.S. Patent No. 6,071,897 issued to DeLuca HF *et al.*). Because of the major drawbacks associated with existing

immunosuppressive modalities, there is a need for a new approach for treating immune disorders, e.g., for inducing immune tolerance in a host.

Thus, in another aspect, the invention provides a method for modulating the activity of an immune cell by contacting the cell with a vitamin D₃ compound of
5 formula I or otherwise described herein.

In one embodiment, the present invention provides a method for suppressing immune activity in an immune cell by contacting a pathological or non-pathological immune cell with an effective amount of a vitamin D₃ compound of the invention to thereby inhibit an immune response relative to the cell in the absence of the treatment.
10 The present method can be performed on cells in culture, e.g., *in vitro* or *ex vivo*, or can be performed on cells present in an animal subject, e.g., as part of an *in vivo* therapeutic protocol. *In vivo* treatment can be carried out on a human or other animal subject.

The vitamin D₃ compounds of the invention can be tested initially *in vitro* for their inhibitory effects on T cell proliferation and secretory activity, as described in
15 Reichel, H. *et al.*, (1987) *Proc. Natl. Acad. Sci. USA* 84:3385-3389; Lemire, J. M. *et al.* (1985) *J. Immunol* 34:2032-2035. Alternatively, the immunosuppressive effects can be tested *in vivo* using the various animal models known in the art and summarized by Bouillon, R. *et al.* (1995) *Endocrine Reviews* 16(2) 232 (Tables 6 and 7). For example, animal models for autoimmune disorders, e.g., lupus, thyroiditis, encephalitis, diabetes
20 and nephritis are described in (Lemire J.M. (1992) *J. Cell Biochem.* 49:26-31; Koizumi T. *et al.* (1985) *Int. Arch. Allergy Appl. Immunol.* 77:396-404; Abe J. *et al.* (1990) *Calcium Regulation and Bone Metabolism* 146-151; Fournier C. *et al.* (1990) *Clin. Immunol Immunopathol.* 54:53-63; Lemire J.M. and Archer D.C. (1991) *J. Clin. Invest.* 87:1103-1107; Lemire, J. M. *et al.*, (1994) *Endocrinology* 135 (6):2818-2821; Inaba M. *et al.* (1992) *Metabolism* 41:631-635; Mathieu C. *et al.* (1992) *Diabetes* 41:1491-1495;
25 Mathieu C. *et al.* (1994) *Diabetologia* 37:552-558; Lillevang S.T. *et al.* (1992) *Clin. Exp. Immunol.* 88:301-306, among others). Models for characterizing immunosuppressive activity during organ transplantation, e.g., skin graft, cardiac graft, islet graft, are described in Jordan S.C. *et al.* (1988) v Herrath D (eds) *Molecular, Cellular and Clinical Endocrinology* 346-347; Veyron P. *et al.* (1993) *Transplant Immunol.* 1:72-76; Jordan S.C. (1988) v Herrath D (eds) *Molecular, Cellular and Clinical Endocrinology* 334-335; Lemire J.M. *et al.* (1992) *Transplantation* 54:762-763; Mathieu C. *et al.* (1994) *Transplant Proc.* 26:3128-3129).

After identifying certain test compounds as effective suppressors of an immune response *in vitro*, these compounds can be used *in vivo* as part of a therapeutic protocol. Accordingly, another aspect of the invention provides a method of suppressing an immune response, comprising administering to a subject a pharmaceutical preparation of
5 a vitamin D₃ compounds of the invention, so as to inhibit immune reactions such as graft rejection, autoimmune disorders and inflammation.

In one embodiment, the invention provides a method for treating a subject for a vitamin D₃ associated state, wherein the vitamin D₃ associated state is an ILT3-associated disorder, by administering to the subject an effective amount of a vitamin D₃
10 compound of the invention. In one embodiment, the the ILT3-associated state is an immune disorder. In certain embodiments, the immune disorder is an autoimmune disorder. In other embodiments, the immune disorder is transplant rejection.

For example, the subject vitamin D₃ compound of the invention can be used to inhibit responses in clinical situations where it is desirable to downmodulate T cell
15 responses. For example, in graft-versus-host disease, cases of transplantation, autoimmune diseases (including, for example, diabetes mellitus, arthritis (including rheumatoid arthritis, juvenile rheumatoid arthritis, osteoarthritis, psoriatic arthritis), multiple sclerosis, encephalomyelitis, diabetes, myasthenia gravis, systemic lupus erythematosus, autoimmune thyroiditis, dermatitis (including atopic dermatitis and
20 eczematous dermatitis), psoriasis, Sjögren's Syndrome, including keratoconjunctivitis sicca secondary to Sjögren's Syndrome, alopecia areata, allergic responses due to arthropod bite reactions, Crohn's disease, aphthous ulcer, iritis, conjunctivitis, keratoconjunctivitis, ulcerative colitis, asthma, allergic asthma, cutaneous lupus erythematosus, scleroderma, vaginitis, proctitis, drug eruptions, leprosy reversal
25 reactions, erythema nodosum leprosum, autoimmune uveitis, allergic encephalomyelitis, acute necrotizing hemorrhagic encephalopathy, idiopathic bilateral progressive sensorineural hearing loss, aplastic anemia, pure red cell anemia, idiopathic thrombocytopenia, polychondritis, Wegener's granulomatosis, chronic active hepatitis, Stevens-Johnson syndrome, idiopathic sprue, lichen planus, Crohn's disease, Graves
30 ophthalmopathy, sarcoidosis, primary biliary cirrhosis, uveitis posterior, and interstitial lung fibrosis). Downmodulation of immune activity will also be desirable in cases of allergy such as, atopic allergy.

Another aspect of the invention provides a method of modulating the expression of an immunoglobulin-like transcript 3 (ILT3) surface molecule in a cell. The method includes contacting the cell with a compound of formula I in an amount effective to modulate the expression of an immunoglobulin-like transcript 3 (ILT3) surface molecule in the cell. In one embodiment, cell is within a subject a subject. In another embodiment the modulation is upregulation of expression. In other embodiment, the modulation is downregulation of expression.

A related aspect of the invention provides a method of treating an ILT3-associated disorder in a subject. The method includes administering to the subject a compound of formula I in an amount effective to modulate the expression of an ILT3 surface molecule, thereby treating the ILT3-associated disorder in the subject.

In certain embodiments, the present invention provides methods and compositions for treating immune disorders, such as, for example, autoimmune disorders and transplant rejections, such as graft versus host disease (GVHD). These embodiments of the invention are based on the discovery that vitamin D₃ compounds of the invention are able to modulate the expression of immunoglobulin-like transcript 3 (ILT3) on cells, e.g., antigen-presenting cells.

Accordingly, another aspect of the invention provides a method for inhibiting transplant rejection in a subject. The method includes administering to the subject a compound of formula I in an amount effective to modulate the expression of an ILT3 surface molecule, thereby inhibiting transplant rejection in the subject. In one embodiment, the transplant is an organ transplant. In another embodiment, the transplant is a pancreatic islet transplant. In yet another embodiment, the transplant is a bone marrow transplant.

As described before, determination of a therapeutically effective immunosuppressive amount can be readily made by the attending clinician, as one skilled in the art, by the use of known techniques and by observing results obtained under analogous circumstances. Compounds which are determined to be effective in animals, e.g., dogs, rodents, may be extrapolated accordingly to humans by those skilled in the art. Starting dose/regimen used in animals can be estimated based on prior studies. For example, doses of vitamin D₃ compounds of the invention to treat autoimmune disorders in rodents can be initially estimated in the range of 0.1 g/kg/day to 1 g/kg/day, administered orally or by injection.

Those skilled in the art will know based upon the data obtained in animal studies, the dosage and route of administration in humans is expected to be similar to that in animals. Exemplary dose ranges to be used in humans are from 0.25 to 10 µg/day, preferably 0.5 to 5 µg/day per adult (U.S. Pat. No. 4,341,774).

5

D. Calcium and Phosphate Homeostasis

The present invention also relates to a method of treating in a subject a disorder characterized by deregulation of calcium metabolism. This method comprises contacting a pathological or non-pathological vitamin D₃ responsive cell with an
10 effective amount of a vitamin D₃ compound of the invention to thereby directly or indirectly modulate calcium and phosphate homeostasis. Techniques for detecting calcium fluctuation *in vivo* or *in vitro* are known in the art.

Exemplary Ca⁺⁺ homeostasis related assays include assays that focus on the intestine where intestinal ⁴⁵Ca²⁺ absorption is determined either 1) *in vivo* (Hibberd
15 K.A. and Norman A.W. (1969) *Biochem. Pharmacol.* 18:2347-2355; Hurwitz S. *et al.* (1967) *J. Nutr.* 91:319-323; Bickle D.D. *et al.* (1984) *Endocrinology* 114:260-267), or 2) *in vitro* with everted duodenal sacs (Schachter D. *et al.* (1961) *Am. J. Physiol* 200:1263-1271), or 3) on the genomic induction of calbindin-D_{28k} in the chick or of calbindin-D_{9k} in the rat (Thomasset M. *et al.* (1981) *FEBS Lett.* 127:13-16; Brehier A.
20 and Thomasset M. (1990) *Endocrinology* 127:580-587). The bone-oriented assays include: 1) assessment of bone resorption as determined via the release of Ca²⁺ from bone *in vivo* (in animals fed a zero Ca²⁺ diet) (Hibberd K.A. and Norman A.W. (1969) *Biochem. Pharmacol.* 18:2347-2355; Hurwitz S. *et al.* (1967) *J. Nutr.* 91:319-323), or from bone explants *in vitro* (Bouillon R. *et al.* (1992) *J. Biol. Chem.* 267:3044-3051), 2)
25 measurement of serum osteocalcin levels [osteocalcin is an osteoblast-specific protein that after its synthesis is largely incorporated into the bone matrix, but partially released into the circulation (or tissue culture medium) and thus represents a good marker of bone formation or turnover] (Bouillon R. *et al.* (1992) *Clin. Chem.* 38:2055-2060), or 3) bone ash content (Norman A.W. and Wong R.G. (1972) *J. Nutr.* 102:1709-1718). Only one
30 kidney-oriented assay has been employed. In this assay, urinary Ca²⁺ excretion is determined (Hartenbower D.L. *et al.* (1977) Walter de Gruyter, Berlin pp 587-589); this assay is dependent upon elevations in the serum Ca²⁺ level and may reflect bone Ca²⁺ mobilizing activity more than renal effects. Finally, there is a "soft tissue calcification"

assay that can be used to detect the consequences of administration of a compound of the invention. In this assay a rat is administered an intraperitoneal dose of $^{45}\text{Ca}^{2+}$, followed by seven daily relative high doses of a compound of the invention; in the event of onset of a severe hypercalcemia, soft tissue calcification can be assessed by
5 determination of the $^{45}\text{Ca}^{2+}$ level. In all these assays, vitamin D₃ compounds of the invention are administered to vitamin D-sufficient or -deficient animals, as a single dose or chronically (depending upon the assay protocol), at an appropriate time interval before the end point of the assay is quantified.

In certain embodiments, vitamin D₃ compounds of the invention can be used to
10 modulate bone metabolism. The language "bone metabolism" is intended to include direct or indirect effects in the formation or degeneration of bone structures, *e.g.*, bone formation, bone resorption, etc., which may ultimately affect the concentrations in serum of calcium and phosphate. This term is also intended to include effects of vitamin D₃ compounds in bone cells, *e.g.* osteoclasts and osteoblasts, that may in turn result in
15 bone formation and degeneration. For example, it is known in the art, that vitamin D₃ compounds exert effects on the bone forming cells, the osteoblasts through genomic and non-genomic pathways (Walters M.R. *et al.* (1982) *J. Biol. Chem.* 257:7481-7484; Jurutka P.W. *et al.* (1993) *Biochemistry* 32:8184-8192; Mellon W.S. and DeLuca H.F. (1980) *J. Biol. Chem.* 255:4081-4086). Similarly, vitamin D₃ compounds are known in
20 the art to support different activities of bone resorbing osteoclasts such as the stimulation of differentiation of monocytes and mononuclear phagocytes into osteoclasts (Abe E. *et al.* (1988) *J. Bone Miner Res.* 3:635-645; Takahashi N. *et al.* (1988) *Endocrinology* 123:1504-1510; Udagawa N. *et al.* (1990) *Proc. Natl. Acad. Sci. USA* 87:7260-7264). Accordingly, vitamin D₃ compounds of the invention that modulate the
25 production of bone cells can influence bone formation and degeneration.

The present invention provides a method for modulating bone cell metabolism by contacting a pathological or a non-pathological bone cell with an effective amount of a vitamin D₃ compound of the invention to thereby modulate bone formation and degeneration. The present method can be performed on cells in culture, *e.g.*, *in vitro* or
30 *ex vivo*, or can be performed in cells present in an animal subject, *e.g.*, cells *in vivo*. Exemplary culture systems that can be used include osteoblast cell lines, *e.g.*, ROS 17/2.8 cell line, monocytes, bone marrow culture system (Suda T. *et al.* (1990) *Med. Res. Rev.* 7:333-366; Suda T. *et al.* (1992) *J. Cell Biochem.* 49:53-58) among others.

Selected compounds can be further tested *in vivo*, for example, animal models of osteopetrosis and in human disease (Shapira F. (1993) *Clin. Orthop.* 294:34-44).

In a preferred embodiment, a method for treating osteoporosis is provided, comprising administering to a subject a pharmaceutical preparation of a vitamin D₃ compound of the invention to thereby ameliorate the condition relative to an untreated subject.

Vitamin D₃ compounds of the invention can be tested in ovariectomized animals, *e.g.*, dogs, rodents, to assess the changes in bone mass and bone formation rates in both normal and estrogen-deficient animals. Clinical trials can be conducted in humans by attending clinicians to determine therapeutically effective amounts of the vitamin D₃ compounds of the invention in preventing and treating osteoporosis.

In other embodiments, therapeutic applications of the vitamin D₃ compounds of the invention include treatment of other diseases characterized by metabolic calcium and phosphate deficiencies. Exemplary of such diseases are the following: osteoporosis, osteodystrophy, osteomalacia, rickets, osteitis fibrosa cystica, renal osteodystrophy, osteosclerosis, anti-convulsant treatment, osteopenia, fibrogenesis-imperfecta ossium, secondary hyperparathyroidism, hypoparathyroidism, hyperparathyroidism, cirrhosis, obstructive jaundice, drug induced metabolism, medullary carcinoma, chronic renal disease, hypophosphatemic VDRR, vitamin D-dependent rickets, sarcoidosis, glucocorticoid antagonism, malabsorption syndrome, steatorrhea, tropical sprue, idiopathic hypercalcemia and milk fever.

E. Hormone Secretion

In yet another aspect, the present invention provides a method for modulating hormone secretion of a vitamin D₃- responsive cell, *e.g.*, an endocrine cell. Hormone secretion includes both genomic and non-genomic activities of vitamin D₃ compounds of the invention that control the transcription and processing responsible for secretion of a given hormone *e.g.*, parathyroid hormone (PTH), calcitonin, insulin, prolactin (PRL) and TRH in a vitamin D₃ responsive cell (Bouillon, R. *et al.* (1995) *Endocrine Reviews* 16(2):235-237).

The present method can be performed on cells in culture, *e.g.* *in vitro* or *ex vivo*, or on cells present in an animal subject, *e.g.*, *in vivo*. Vitamin D₃ compounds of the invention can be initially tested *in vitro* using primary cultures of parathyroid cells.

- Other systems that can be used include the testing by prolactin secretion in rat pituitary tumor cells, e.g., GH4C1 cell line (Wark J.D. and Tashjian Jr. A.H. (1982) *Endocrinology* 111:1755-1757; Wark J. D. and Tashjian Jr. A.H. (1983) *J. Biol. Chem.* 258:2118-2121; Wark J.D. and Gurtler V. (1986) *Biochem. J.* 233:513-518) and TRH secretion in GH4C1 cells. Alternatively, the effects of vitamin D₃ compounds of the invention can be characterized *in vivo* using animals models as described in Nko M. *et al.* (1982) *Miner Electrolyte Metab.* 5:67-75; Oberg F. *et al.* (1993) *J. Immunol.* 150:3487-3495; Bar-Shavit Z. *et al.* (1986) *Endocrinology* 118:679-686; Testa U. *et al.* (1993) *J. Immunol.* 150:2418-2430; Nakamaki T. *et al.* (1992) *Anticancer Res.* 12:1331-1337; Weinberg J.B. and Larrick J.W. (1987) *Blood* 70:994-1002; Chambaut-Guérin A.M. and Thomopoulos P. (1991) *Eur. Cytokine New.* 2:355; Yoshida M. *et al.* (1992) *Anticancer Res.* 12:1947-1952; Momparler R.L. *et al.* (1993) *Leukemia* 7:17-20; Eisman J.A. (1994) *Kanis JA (eds) Bone and Mineral Research* 2:45-76; Veyron P. *et al.* (1993) *Transplant Immunol.* 1:72-76; Gross M. *et al.* (1986) *J Bone Miner Res.* 1:457-467; Costa E.M. *et al.* (1985) *Endocrinology* 117:2203-2210; Koga M. *et al.* (1988) *Cancer Res.* 48:2734-2739; Franceschi R.T. *et al.* (1994) *J. Cell Physiol.* 123:401-409; Cross H.S. *et al.* (1993) *Naunyn Schmiedeberg's Arch. Pharmacol.* 347:105-110; Zhao X. and Feldman D. (1993) *Endocrinology* 132:1808-1814; Skowronski R.J. *et al.* (1993) *Endocrinology* 132:1952-1960; Henry H.L. and Norman A.W. (1975) *Biochem. Biophys. Res. Commun.* 62:781-788; Weckler W.R. *et al.* (1980) *Arch. Biochem. Biophys.* 201:95-103; Brumbaugh P.F. *et al.* (1975) *Am. J. Physiol.* 238:384-388; Oldham S.B. *et al.* (1979) *Endocrinology* 104:248-254; Chertow B.S. *et al.* (1975) *J. Clin Invest.* 56:668-678; Canterbury J.M. *et al.* (1978) *J. Clin. Invest.* 61:1375-1383; Quesad J.M. *et al.* (1992) *J. Clin. Endocrinol. Metab.* 75:494-501.
- 25 In certain embodiments, the vitamin D₃ compounds of the present invention can be used to inhibit parathyroid hormone (PTH) processing, e.g., transcriptional, translational processing, and/or secretion of a parathyroid cell as part of a therapeutic protocol. Therapeutic methods using these compounds can be readily applied to all diseases, involving direct or indirect effects of PTH activity, e.g., primary or secondary responses.

Accordingly, therapeutic applications for the vitamin D₃ compounds of the invention include treating diseases such as secondary hyperparathyroidism of chronic renal failure (Slatopolsky E. *et al.* (1990) *Kidney Int.* 38:S41-S47; Brown A.J. *et al.*

(1989) *J. Clin. Invest.* 84:728-732). Determination of therapeutically affective amounts and dose regimen can be performed by the skilled artisan using the data described in the art.

5 F. Protection Against Neuronal Loss

In yet another aspect, the present invention provides a method of protecting against neuronal loss by contacting a vitamin D₃ responsive cell, e.g., a neuronal cell, with a vitamin D₃ compound of the invention to prevent or retard neuron loss. The language "protecting against" is intended to include prevention, retardation, and/or

10 termination of deterioration, impairment, or death of a neurons.

Neuron loss can be the result of any condition of a neuron in which its normal function is compromised. Neuron deterioration can be the result of any condition which compromises neuron function which is likely to lead to neuron loss. Neuron function can be compromised by, for example, altered biochemistry, physiology, or anatomy of a
15 neuron. Deterioration of a neuron may include membrane, dendritic, or synaptic changes which are detrimental to normal neuronal functioning. The cause of the neuron deterioration, impairment, and/or death may be unknown. Alternatively, it may be the result of age- and/or disease-related changes which occur in the nervous system of a subject.

20 When neuron loss is described herein as "age-related", it is intended to include neuron loss resulting from known and unknown bodily changes of a subject which are associated with aging. When neuron loss is described herein as "disease-related", it is intended to include neuron loss resulting from known and unknown bodily changes of a subject which are associated with disease. It should be understood, however, that these
25 terms are not mutually exclusive and that, in fact, many conditions that result in the loss of neurons are both age- and disease-related.

Exemplary age-related diseases associated with neuron loss and changes in neuronal morphology include, for example, Alzheimer's Disease, Pick's Disease, Parkinson's Disease, Vascular Disease, Huntington's Disease, and Age-Associated
30 Memory Impairment. In Alzheimer's Disease patients, neuron loss is most notable in the hippocampus, frontal, parietal, and anterior temporal cortices, amygdala, and the olfactory system. The most prominently affected zones of the hippocampus include the CA1 region, the subiculum, and the entorhinal cortex. Memory loss is considered the

earliest and most representative cognitive change because the hippocampus is well known to play a crucial role in memory. Pick's Disease is characterized by severe neuronal degeneration in the neocortex of the frontal and anterior temporal lobes which is sometimes accompanied by death of neurons in the striatum. Parkinson's Disease can be identified by the loss of neurons in the substantia nigra and the locus ceruleus. Huntington's Disease is characterized by degeneration of the intrastriatal and cortical cholinergic neurons and GABA-ergic neurons. Parkinson's and Huntington's Diseases are usually associated with movement disorders, but often show cognitive impairment (memory loss) as well.

10 Age-Associated Memory Impairment (AAMI) is another age-associated disorder that is characterized by memory loss in healthy, elderly individuals in the later decades of life. Crook, T. *et al.* (1986) *Devel. Neuropsych.* 2(4):261-276. Presently, the neural basis for AAMI has not been precisely defined. However, neuron death with aging has been reported to occur in many species in brain regions implicated in memory, including
15 cortex, hippocampus, amygdala, basal ganglia, cholinergic basal forebrain, locus ceruleus, raphe nuclei, and cerebellum. Crook, T. *et al.* (1986) *Devel. Neuropsych.* 2(4):261-276.

Vitamin D₃ compounds of the invention can protect against neuron loss by genomic or non-genomic mechanisms. Nuclear vitamin D₃ receptors are well known to
20 exist in the periphery but have also been found in the brain, particularly in the hippocampus and neocortex. Non-genomic mechanisms may also prevent or retard neuron loss by regulating intraneuronal and/or peripheral calcium and phosphate levels. Furthermore, vitamin D₃ compounds of the invention may protect against neuronal loss by acting indirectly, *e.g.*, by modulating serum PTH levels. For example, a positive
25 correlation has been demonstrated between serum PTH levels and cognitive decline in Alzheimer's Disease.

The present method can be performed on cells in culture, *e.g. in vitro* or *ex vivo*, or on cells present in an animal subject, *e.g., in vivo*. Vitamin D₃ compounds of the invention can be initially tested *in vitro* using neurons from embryonic rodent pups (*See*
30 *e.g.* U.S. Patent No. 5,179,109-fetal rat tissue culture), or other mammalian (*See e.g.* U.S. Patent No. 5,089,517-fetal mouse tissue culture) or non-mammalian animal models. These culture systems have been used to characterize the protection of peripheral, as well as, central nervous system neurons in animal or tissue culture models

of ischemia, stroke, trauma, nerve crush, Alzheimer's Disease, Pick's Disease, and Parkinson's Disease, among others. Examples of *in vitro* systems to study the prevention of destruction of neocortical neurons include using *in vitro* cultures of fetal mouse neurons and glial cells previously exposed to various glutamate agonists, such as kainate, NMDA, and α -amino-3-hydroxy-5-methyl-4-isoxazoleprionate (AMPA). U.S. Patent No. 5,089,517. *See also* U.S. Patent No. 5,170,109 (treatment of rat cortical/hippocampal neuron cultures with glutamate prior to treatment with neuroprotective compound); U.S. Patent Nos. 5,163,196 and 5,196,421 (neuroprotective excitatory amino acid receptor antagonists inhibit glycine, kainate, AMPA receptor binding in rats).

Alternatively, the effects of vitamin D₃ compounds of the invention can be characterized *in vivo* using animals models. Neuron deterioration in these model systems is often induced by experimental trauma or intervention (*e.g.* application of toxins, nerve crush, interruption of oxygen supply).

G. Smooth Muscle Cells

In yet another aspect, the present invention provides a method of modulating the activity of a vascular smooth muscle cell by contacting a vitamin D₃-responsive smooth muscle cell with a vitamin D₃ compound of the invention to activate or, preferably, inhibit the activity of the cell. The language "activity of a smooth muscle cell" is intended to include any activity of a smooth muscle cell, such as proliferation, migration, adhesion and/or metabolism.

In certain embodiments, the vitamin D₃ compounds of the invention can be used to treat diseases and conditions associated with aberrant activity of a vitamin D₃-responsive smooth muscle cell. For example, the present invention can be used in the treatment of hyperproliferative vascular diseases, such as hypertension induced vascular remodeling, vascular restenosis and atherosclerosis. In other embodiments, the compounds of the present invention can be used in treating disorders characterized by aberrant metabolism of a vitamin D₃-responsive smooth muscle cell, *e.g.*, arterial hypertension.

The present method can be performed on cells in culture, *e.g.* *in vitro* or *ex vivo*, or on cells present in an animal subject, *e.g.*, *in vivo*. Vitamin D₃ compounds of the

invention can be initially tested *in vitro* as described in Catellot *et al.* (1982), *J. Biol. Chem.* 257(19): 11256.

H. Suppression of Renin Expression and Treatment of Hypertension

5 The compounds of the present invention control blood pressure by the suppression of rennin expression and are useful as antihypertensive agents. Renin-angiotensin regulatory cascade plays a significant role in the regulation of blood pressure, electrolyte and volume homeostasis (Y.C. Li, Abstract, *DeLuca Symposium on Vitamin D₃*, Taos, New Mexico, June 15 - June 19, 2002, p. 18). Thus, the invention
10 provides a method of treating a subject for a vitamin D₃ associated state, wherein the vitamin D₃ associated state is a disorder characterized by an aberrant activity of a cell that expresses renin. The method includes administering to the subject an effective amount of a compound of formula I, such that renin expression by the cell is suppressed, and the subject is thereby treated for hypertension.

15

I. Treatment of Urogenital Disorders

 The invention also provides a method for for treating a subject for a urogenital disorder. The method comprises administering to the subject an effective amount of a vitamin D₃ compound of formula I above, such that the subject is treated for the
20 urogenital disorder.

 In one embodiment, the urogenital disorder comprises bladder dysfunction, especially bladder dysfunction related to morphological bladder changes. The term bladder dysfunction as used in this embodiment does not include cancer of the bladder and associated urogenital organs.

25 Morphological bladder changes, including a progressive de-nervation and hypertrophy of the bladder wall are frequent histological findings in patients with different bladder disorders such as overactive bladder and clinical BPH. The increase in tension and/or strain on the bladder observed in these conditions has been shown to be associated with cellular and molecular alterations, *e.g.*, in cytoskeletal and contractile
30 proteins, in mitochondrial function, and in various enzyme activities of the smooth muscle cells. The growth of the bladder wall also involves alterations in its extracellular matrix and non-smooth muscle components.

 These changes in the bladder are associated with the storage (irritative)

symptoms, in particular frequency, urgency and nocturia. These symptoms affect the social, psychological, domestic, occupational, physical and sexual lives of the patients leading to a profound, negative impact on their quality of life.

Also included within urogenital disorders is benign prostatic hyperplasia (BPH). Thus the invention also provides a method for treatment of BPH comprising administering to a subject an effective amount of a vitamin D₃ compound of formula I above, such that the subject is treated for BPH.

BPH is commonly associated with enlargement of the gland (prostate) leading to bladder outlet obstruction (BOO) and symptoms secondary to BOO. However, BPH is also associated with morphological bladder changes, including a progressive denervation and hypertrophy of the bladder wall, the latter possibly as a consequence of increased functional demands. Thus, the compounds of the invention are useful for the treatment of storage (irritative) symptoms of BPH, as well as for bladder outlet obstruction caused by BPH.

Urogenital disorders in accordance with the invention also include interstitial cystitis. Thus, in another embodiment, the invention also provides a method for treatment of interstitial cystitis comprising administering to a subject an effective amount of a vitamin D₃ compound of formula I above, such that the subject is treated for interstitial cystitis.

Interstitial cystitis (IC) is a chronic inflammatory bladder disease characterized by pelvic pain, urinary urgency and frequency. Unlike other bladder dysfunction conditions, IC is characterized by chronic inflammation of the bladder wall which is responsible for the symptomatology. In other words, the cause of the abnormal bladder contractility is the chronic inflammation and as a consequence the treatment should target this etiological component. In fact, the traditional treatment of bladder dysfunctions, like overactive bladder, with smooth muscle relaxant agents, is not effective in patients with IC.

4. PHARMACEUTICAL COMPOSITIONS

The invention also provides a pharmaceutical composition, comprising an effective amount of a vitamin D₃ compound of formula I or otherwise described herein and a pharmaceutically acceptable carrier. In a further embodiment, the effective amount is effective to treat a vitamin D₃ associated state, as described previously.

In an embodiment, the vitamin D₃ compound is administered to the subject using a pharmaceutically-acceptable formulation, *e.g.*, a pharmaceutically-acceptable formulation that provides sustained delivery of the vitamin D₃ compound to a subject for at least 12 hours, 24 hours, 36 hours, 48 hours, one week, two weeks, three weeks, or
5 four weeks after the pharmaceutically-acceptable formulation is administered to the subject.

In certain embodiments, these pharmaceutical compositions are suitable for topical or oral administration to a subject. In other embodiments, as described in detail below, the pharmaceutical compositions of the present invention may be specially
10 formulated for administration in solid or liquid form, including those adapted for the following: (1) oral administration, for example, drenches (aqueous or non-aqueous solutions or suspensions), tablets, boluses, powders, granules, pastes; (2) parenteral administration, for example, by subcutaneous, intramuscular or intravenous injection as, for example, a sterile solution or suspension; (3) topical application, for example, as a
15 cream, ointment or spray applied to the skin; (4) intravaginally or intrarectally, for example, as a pessary, cream or foam; or (5) aerosol, for example, as an aqueous aerosol, liposomal preparation or solid particles containing the compound.

The phrase "pharmaceutically acceptable" refers to those vitamin D₃ compounds of the present invention, compositions containing such compounds, and/or dosage forms
20 which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of human beings and animals without excessive toxicity, irritation, allergic response, or other problem or complication, commensurate with a reasonable benefit/risk ratio.

The phrase "pharmaceutically-acceptable carrier" includes pharmaceutically-
25 acceptable material, composition or vehicle, such as a liquid or solid filler, diluent, excipient, solvent or encapsulating material, involved in carrying or transporting the subject chemical from one organ, or portion of the body, to another organ, or portion of the body. Each carrier must be "acceptable" in the sense of being compatible with the other ingredients of the formulation and not injurious to the patient. Some examples of
30 materials which can serve as pharmaceutically-acceptable carriers include: (1) sugars, such as lactose, glucose and sucrose; (2) starches, such as corn starch and potato starch; (3) cellulose, and its derivatives, such as sodium carboxymethyl cellulose, ethyl cellulose and cellulose acetate; (4) powdered tragacanth; (5) malt; (6) gelatin; (7) talc;

(8) excipients, such as cocoa butter and suppository waxes; (9) oils, such as peanut oil, cottonseed oil, safflower oil, sesame oil, olive oil, corn oil and soybean oil; (10) glycols, such as propylene glycol; (11) polyols, such as glycerin, sorbitol, mannitol and polyethylene glycol; (12) esters, such as ethyl oleate and ethyl laurate; (13) agar; (14) buffering agents, such as magnesium hydroxide and aluminum hydroxide; (15) alginic acid; (16) pyrogen-free water; (17) isotonic saline; (18) Ringer's solution; (19) ethyl alcohol; (20) phosphate buffer solutions; and (21) other non-toxic compatible substances employed in pharmaceutical formulations.

Wetting agents, emulsifiers and lubricants, such as sodium lauryl sulfate and magnesium stearate, as well as coloring agents, release agents, coating agents, sweetening, flavoring and perfuming agents, preservatives and antioxidants can also be present in the compositions.

Examples of pharmaceutically-acceptable antioxidants include: (1) water soluble antioxidants, such as ascorbic acid, cysteine hydrochloride, sodium bisulfate, sodium metabisulfite, sodium sulfite and the like; (2) oil-soluble antioxidants, such as ascorbyl palmitate, butylated hydroxyanisole (BHA), butylated hydroxytoluene (BHT), lecithin, propyl gallate, alpha-tocopherol, and the like; and (3) metal chelating agents, such as citric acid, ethylenediamine tetraacetic acid (EDTA), sorbitol, tartaric acid, phosphoric acid, and the like.

Compositions containing a vitamin D₃ compound(s) include those suitable for oral, nasal, topical (including buccal and sublingual), rectal, vaginal, aerosol and/or parenteral administration. The compositions may conveniently be presented in unit dosage form and may be prepared by any methods well known in the art of pharmacy. The amount of active ingredient which can be combined with a carrier material to produce a single dosage form will vary depending upon the host being treated, the particular mode of administration. The amount of active ingredient which can be combined with a carrier material to produce a single dosage form will generally be that amount of the compound which produces a therapeutic effect. Generally, out of one hundred per cent, this amount will range from about 1 per cent to about ninety-nine percent of active ingredient, preferably from about 5 per cent to about 70 per cent, most preferably from about 10 per cent to about 30 per cent.

Methods of preparing these compositions include the step of bringing into association a vitamin D₃ compound(s) with the carrier and, optionally, one or more

accessory ingredients. In general, the formulations are prepared by uniformly and intimately bringing into association a vitamin D₃ compound with liquid carriers, or finely divided solid carriers, or both, and then, if necessary, shaping the product.

Compositions of the invention suitable for oral administration may be in the form of capsules, cachets, pills, tablets, lozenges (using a flavored basis, usually sucrose and acacia or tragacanth), powders, granules, or as a solution or a suspension in an aqueous or non-aqueous liquid, or as an oil-in-water or water-in-oil liquid emulsion, or as an elixir or syrup, or as pastilles (using an inert base, such as gelatin and glycerin, or sucrose and acacia) and/or as mouth washes and the like, each containing a predetermined amount of a vitamin D₃ compound(s) as an active ingredient. A compound may also be administered as a bolus, electuary or paste.

In solid dosage forms of the invention for oral administration (capsules, tablets, pills, dragees, powders, granules and the like), the active ingredient is mixed with one or more pharmaceutically-acceptable carriers, such as sodium citrate or dicalcium phosphate, and/or any of the following: (1) fillers or extenders, such as starches, lactose, sucrose, glucose, mannitol, and/or silicic acid; (2) binders, such as, for example, carboxymethylcellulose, alginates, gelatin, polyvinyl pyrrolidone, sucrose and/or acacia; (3) humectants, such as glycerol; (4) disintegrating agents, such as agar-agar, calcium carbonate, potato or tapioca starch, alginic acid, certain silicates, and sodium carbonate; (5) solution retarding agents, such as paraffin; (6) absorption accelerators, such as quaternary ammonium compounds; (7) wetting agents, such as, for example, acetyl alcohol and glycerol monostearate; (8) absorbents, such as kaolin and bentonite clay; (9) lubricants, such as talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate, and mixtures thereof; and (10) coloring agents. In the case of capsules, tablets and pills, the pharmaceutical compositions may also comprise buffering agents. Solid compositions of a similar type may also be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugars, as well as high molecular weight polyethylene glycols and the like.

A tablet may be made by compression or molding, optionally with one or more accessory ingredients. Compressed tablets may be prepared using binder (for example, gelatin or hydroxypropylmethyl cellulose), lubricant, inert diluent, preservative, disintegrant (for example, sodium starch glycolate or cross-linked sodium carboxymethyl cellulose), surface-active or dispersing agent. Molded tablets may be

made by molding in a suitable machine a mixture of the powdered active ingredient moistened with an inert liquid diluent.

The tablets, and other solid dosage forms of the pharmaceutical compositions of the present invention, such as dragees, capsules, pills and granules, may optionally be scored or prepared with coatings and shells, such as enteric coatings and other coatings well known in the pharmaceutical-formulating art. They may also be formulated so as to provide slow or controlled release of the active ingredient therein using, for example, hydroxypropylmethyl cellulose in varying proportions to provide the desired release profile, other polymer matrices, liposomes and/or microspheres. They may be sterilized by, for example, filtration through a bacteria-retaining filter, or by incorporating sterilizing agents in the form of sterile solid compositions which can be dissolved in sterile water, or some other sterile injectable medium immediately before use. These compositions may also optionally contain opacifying agents and may be of a composition that they release the active ingredient(s) only, or preferentially, in a certain portion of the gastrointestinal tract, optionally, in a delayed manner. Examples of embedding compositions which can be used include polymeric substances and waxes. The active ingredient can also be in micro-encapsulated form, if appropriate, with one or more of the above-described excipients.

Liquid dosage forms for oral administration of the vitamin D₃ compound(s) include pharmaceutically-acceptable emulsions, microemulsions, solutions, suspensions, syrups and elixirs. In addition to the active ingredient, the liquid dosage forms may contain inert diluents commonly used in the art, such as, for example, water or other solvents, solubilizing agents and emulsifiers, such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, oils (in particular, cottonseed, groundnut, corn, germ, olive, castor and sesame oils), glycerol, tetrahydrofuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan, and mixtures thereof.

In addition to inert diluents, the oral compositions can include adjuvants such as wetting agents, emulsifying and suspending agents, sweetening, flavoring, coloring, perfuming and preservative agents.

Suspensions, in addition to the active vitamin D₃ compound(s) may contain suspending agents as, for example, ethoxylated isostearyl alcohols, polyoxyethylene

sorbitol and sorbitan esters, microcrystalline cellulose, aluminum metahydroxide, bentonite, agar-agar and tragacanth, and mixtures thereof.

Pharmaceutical compositions of the invention for rectal or vaginal administration may be presented as a suppository, which may be prepared by mixing one or more
5 vitamin D₃ compound(s) with one or more suitable nonirritating excipients or carriers comprising, for example, cocoa butter, polyethylene glycol, a suppository wax or a salicylate, and which is solid at room temperature, but liquid at body temperature and, therefore, will melt in the rectum or vaginal cavity and release the active agent.

Compositions of the present invention which are suitable for vaginal
10 administration also include pessaries, tampons, creams, gels, pastes, foams or spray formulations containing such carriers as are known in the art to be appropriate.

Dosage forms for the topical or transdermal administration of a vitamin D₃ compound(s) include powders, sprays, ointments, pastes, creams, lotions, gels, solutions, patches and inhalants. The active vitamin D₃ compound(s) may be mixed under sterile
15 conditions with a pharmaceutically-acceptable carrier, and with any preservatives, buffers, or propellants which may be required.

The ointments, pastes, creams and gels may contain, in addition to vitamin D₃ compound(s) of the present invention, excipients, such as animal and vegetable fats, oils, waxes, paraffins, starch, tragacanth, cellulose derivatives, polyethylene glycols,
20 silicones, bentonites, silicic acid, talc and zinc oxide, or mixtures thereof.

Powders and sprays can contain, in addition to a vitamin D₃ compound(s), excipients such as lactose, talc, silicic acid, aluminum hydroxide, calcium silicates and polyamide powder, or mixtures of these substances. Sprays can additionally contain customary propellants, such as chlorofluorohydrocarbons and volatile unsubstituted
25 hydrocarbons, such as butane and propane.

The vitamin D₃ compound(s) can be alternatively administered by aerosol. This is accomplished by preparing an aqueous aerosol, liposomal preparation or solid particles containing the compound. A nonaqueous (*e.g.*, fluorocarbon propellant) suspension could be used. Sonic nebulizers are preferred because they minimize
30 exposing the agent to shear, which can result in degradation of the compound.

Ordinarily, an aqueous aerosol is made by formulating an aqueous solution or suspension of the agent together with conventional pharmaceutically-acceptable carriers and stabilizers. The carriers and stabilizers vary with the requirements of the particular

compound, but typically include nonionic surfactants (Tweens, Pluronics, or polyethylene glycol), innocuous proteins like serum albumin, sorbitan esters, oleic acid, lecithin, amino acids such as glycine, buffers, salts, sugars or sugar alcohols. Aerosols generally are prepared from isotonic solutions.

- 5 Transdermal patches have the added advantage of providing controlled delivery of a vitamin D₃ compound(s) to the body. Such dosage forms can be made by dissolving or dispersing the agent in the proper medium. Absorption enhancers can also be used to increase the flux of the active ingredient across the skin. The rate of such flux can be controlled by either providing a rate controlling membrane or dispersing the active
10 ingredient in a polymer matrix or gel.

Ophthalmic formulations, eye ointments, powders, solutions and the like, are also contemplated as being within the scope of the invention.

- Pharmaceutical compositions of the invention suitable for parenteral administration comprise one or more vitamin D₃ compound(s) in combination with one
15 or more pharmaceutically-acceptable sterile isotonic aqueous or nonaqueous solutions, dispersions, suspensions or emulsions, or sterile powders which may be reconstituted into sterile injectable solutions or dispersions just prior to use, which may contain antioxidants, buffers, bacteriostats, solutes which render the formulation isotonic with the blood of the intended recipient or suspending or thickening agents.

- 20 Examples of suitable aqueous and nonaqueous carriers which may be employed in the pharmaceutical compositions of the invention include water, ethanol, polyols (such as glycerol, propylene glycol, polyethylene glycol, and the like), and suitable mixtures thereof, vegetable oils, such as olive oil, and injectable organic esters, such as ethyl oleate. Proper fluidity can be maintained, for example, by the use of coating
25 materials, such as lecithin, by the maintenance of the required particle size in the case of dispersions, and by the use of surfactants.

- These compositions may also contain adjuvants such as preservatives, wetting agents, emulsifying agents and dispersing agents. Prevention of the action of microorganisms may be ensured by the inclusion of various antibacterial and antifungal
30 agents, for example, paraben, chlorobutanol, phenol sorbic acid, and the like. It may also be desirable to include isotonic agents, such as sugars, sodium chloride, and the like into the compositions. In addition, prolonged absorption of the injectable pharmaceutical

form may be brought about by the inclusion of agents which delay absorption such as aluminum monostearate and gelatin.

In some cases, in order to prolong the effect of a drug, it is desirable to slow the absorption of the drug from subcutaneous or intramuscular injection. This may be accomplished by the use of a liquid suspension of crystalline or amorphous material having poor water solubility. The rate of absorption of the drug then depends upon its rate of dissolution which, in turn, may depend upon crystal size and crystalline form. Alternatively, delayed absorption of a parenterally-administered drug form is accomplished by dissolving or suspending the drug in an oil vehicle.

Injectable depot forms are made by forming microencapsule matrices of vitamin D₃ compound(s) in biodegradable polymers such as polylactide-polyglycolide. Depending on the ratio of drug to polymer, and the nature of the particular polymer employed, the rate of drug release can be controlled. Examples of other biodegradable polymers include poly(orthoesters) and poly(anhydrides). Depot injectable formulations are also prepared by entrapping the drug in liposomes or microemulsions which are compatible with body tissue.

When the vitamin D₃ compound(s) are administered as pharmaceuticals, to humans and animals, they can be given per se or as a pharmaceutical composition containing, for example, 0.1 to 99.5% (more preferably, 0.5 to 90%) of active ingredient in combination with a pharmaceutically-acceptable carrier.

Regardless of the route of administration selected, the vitamin D₃ compound(s), which may be used in a suitable hydrated form, and/or the pharmaceutical compositions of the present invention, are formulated into pharmaceutically-acceptable dosage forms by conventional methods known to those of skill in the art.

Actual dosage levels and time course of administration of the active ingredients in the pharmaceutical compositions of the invention may be varied so as to obtain an amount of the active ingredient which is effective to achieve the desired therapeutic response for a particular patient, composition, and mode of administration, without being toxic to the patient. An exemplary dose range is from 0.1 to 10 mg per day.


A preferred dose of the vitamin D₃ compound for the present invention is the maximum that a patient can tolerate and not develop serious hypercalcemia. Preferably, the vitamin D₃ compound of the present invention is administered at a concentration of about 0.001 µg to about 100 µg per kilogram of body weight, about 0.001 – about 10

$\mu\text{g/kg}$ or about $0.001 \mu\text{g}$ – about $100 \mu\text{g/kg}$ of body weight. Ranges intermediate to the above-recited values are also intended to be part of the invention.

5. SYNTHESIS OF COMPOUNDS OF THE INVENTION

- 5 Compounds of the invention can be synthesized by methods described in this section, the examples, and the chemical literature. In particular, compounds of formula I of the invention are prepared as shown in Scheme 1 below. Accordingly, compounds of formula I are prepared by coupling compounds of formula II with compounds of formula III in tetrahydrofuran with *n*-butyllithium as a base to give compounds of formula IV. Subsequent removal of the protecting silyl groups ($R_1 = \text{OSi}(\text{CH}_3)_2\text{t-Bu}$) affords the 1,3 dihydroxy vitamin D₃ compound of formula I ($R_1 = \text{OH}$, $R_2 = \text{OH}$). Acylation at the 1 and/or 3 positions is achieved using methods well-known in the art. For example, preparation of the 1,3 diacetox compounds of formula I ($R_1 = R_2 = \text{OAc}$) requires additional acetylation with acetic anhydride and pyridine, as shown in Scheme 2 and described in Example 1.
- 10
- 15


Scheme 1



wherein X_1 , X_2 , R_3 , R_4 , R_5 and R_6 are as defined above.

Scheme 2

5




Referring to Schemes 1 and 3, compounds of formula II are known compounds, and are prepared starting from the known epoxy-ketone of formula V. The compound of formula V is converted to the epoxy-olefin of formula VII by a Wittig reaction.

- 10 Reduction with LiAlH_4 to the compound VIII and protection of the hydroxy group resulted in compound IX. Then, the ene reaction of formula IX with the known hydroxy-conjugated ketone X ($R_5 = R_6 = \text{CH}_3$) in tetrahydrofuran, in the presence of Lewis acid $(\text{CH}_3)_2\text{AlCl}$, provides the compound XI featuring the C,D-rings and full side chain of the target vitamin D analogs. Finally, removal of the silyl group and
- 15 oxidation provides the key intermediate, Ketone of formula III.

20

Scheme 3



EXEMPLIFICATION OF THE INVENTION

The invention is further illustrated by the following examples which should in no way should be construed as being further limiting.

5 Synthesis of Compounds of the Invention

Experimental

All operations involving vitamin D₃ analogs were conducted in amber-colored glassware in a nitrogen atmosphere. Tetrahydrofuran was distilled from sodium-benzophenone ketyl just prior to its use and solutions of solutes were dried with sodium
10 sulfate. Melting points were determined on a Thomas-Hoover capillary apparatus and are uncorrected. Optical rotations were measured at 25 °C. ¹H NMR spectra were recorded at 400 MHz in CDCl₃ unless indicated otherwise. TLC was carried out on silica gel plates (Merck PF-254) with visualization under short-wavelength UV light or by spraying the plates with 10% phosphomolybdic acid in methanol followed by
15 heating. Flash chromatography was carried out on 40-65 µm mesh silica gel. Preparative HPLC was performed on a 5×50 cm column and 15-30 µm mesh silica gel at a flow rate of 100 ml/min.

EXAMPLE 1

20 *Synthesis of 1,3-O-Diacetyl-1,25-Dihydroxy-16-ene-24-Keto-19-nor-Cholecalciferol (2)*

Referring to Scheme 2, 0.032 g of 1,25-dihydroxy-16-ene-24-keto-19-nor-cholecalciferol (1) was dissolved in 0.8 ml pyridine, cooled in bath and treated with 0.2
25 ml acetic anhydride for 7 hours at room temperature and for 14 hours in a refrigerator. It was then diluted with 1 ml of water, stirred for 10 min in an ice bath, diluted with 5 ml water and 20 ml ethyl acetate. The organic layer was washed with 3 x 5 ml of water, then with 5 ml saturated sodium bicarbonate, then with brine, dried over sodium sulfate and evaporated. The oily residue was taken up in 1:6 ethyl acetate-hexane, then flash
30 chromatographed on a 13.5 x 110 mm column using 1:6 ethyl acetate-hexane as mobile phase for fractions 1 – 5, 1:4 ethyl acetate-hexane for the remaining fractions. Fractions 11 – 14 were pooled and evaporated to give 0.0184 g of the title compound (2).

EXAMPLE 2***Determination of Maximum Tolerated Dose (MTD) of 1,3-O-Diacetyl-1,25-Dihydroxy-16-ene-24-Keto-19-nor-Cholecalciferol (2)***

5

The maximum tolerated dose of the vitamin D₃ compounds of the invention were determined in eight week-old female C57BL/6 mice (3 mice/group) dosed orally (0.1 ml/mouse) with various concentrations of Vitamin D₃ analogs daily for four days. Analogs were formulated in miglyol for a final concentration of 0.01, 0.03, 0.1 0.3, 1, 3, 10, 30, 100
10 and 300 µg/kg when given at 0.1 ml/mouse p.o. daily. Blood for serum calcium assay was drawn by tail bleed on day five, the final day of the study. Serum calcium levels were determined using a colorimetric assay (Sigma Diagnostics, procedure no. 597). The highest dose of analog tolerated without inducing hypercalcemia (serum calcium >10.7 mg/dl) was taken as the maximum tolerated dose (MTD). Table 1 shows the relative MTD
15 for compound (2) and compound (1). Notably, compound (2) has an MTD that is more than 300 times greater than compound (1).

EXAMPLE 3***Immunological Assay of 1,3-O-Diacetyl-1,25-Dihydroxy-16-ene-24-Keto-19-nor-Cholecalciferol (2)***

20

Immature dendritic cells (DC) were prepared as described in Romani, N. *et al.* (Romani, N. *et al.* (1996) *J. Immunol. Meth.* 196:137). IFN-γ production by allogeneic T cell activation in the mixed leukocyte response (MLR) was determined as described in
25 Penna, G., *et al.*, *J Immunol.*, 164: 2405-2411 (2000).

Briefly, peripheral blood mononuclear cells (PBMC) were separated from buffy coats by Ficoll gradient and the same number (3x10⁵) of allogeneic PBMC from 2 different donors were co-cultured in 96-well flat-bottom plates. After 5 days, IFN-γ production in the MLR assay was measured by ELISA and the results expressed as amount (nM) of test
30 compound required to induce 50% inhibition of IFN-γ production (IC₅₀). The results are summarised in Table 1.

Table 1

Compound	MTD (mice) μg/kg	INF-γ IC ₅₀ pM
<i>1,25(OH)₂D₃</i>	1	22
<i>1,25 Dihydroxy-16-ene-24-Keto-19-nor Cholecalciferol (1)</i>	1	29
<i>1,3-O-Diacetyl-1,25 Dihydroxy-16-ene-24-Keto-19-nor Cholecalciferol (2)</i>	>300	8

EXAMPLE 4

5 ***Immunological Assay For Modulation of ILT3 Activity of 1,3-O-Diacetyl-1,25-Dihydroxy-16-ene-24-Keto-19-nor-Cholecalciferol (2)***

10 **I. Methods**

Immature dendritic cells (DC) were prepared as described in Romani, N. *et al.* (Romani, N. *et al.* (1996) J. Immunol. Meth. 196:137). Briefly, peripheral blood mononuclear cells (PBMCs) were obtained from a buffy coat by ficoll-hypaque gradient (Pharmacia Biotec AB, Upsala, Sweden) and monocytes isolated from PBMCs by negative selected with a monocyte isolation kit (Milteny, Biotech, Bergish Gladblach, Germany). After isolation, monocytes were cultured for 6 to 7 days at a cell density of $1-2 \times 10^6$ in RPMI medium supplemented with 5% FetalClone I (Hyclone Laboratories, Logan, Utah) 2 mM L-glutamine, 50 mg/ml gentamicin, 1 mM sodium pyruvate and 1% nonessential amino acids, containing 800 U/ml GM-CSF (Mielogen 300, Schering-Plough) and 10 ng/ml interleukin (IL)-4 (PharMingen, San Diego, California). Every other day, approximately 20% of the medium was removed and replaced by the same volume of fresh medium containing GM-CSF and IL-4.

After 6 to 7 days of culture, non-adherent cells (representing immature Dendritic Cells) were harvested and cultured for 24 hours in the presence of graded doses of the following vitamin D compounds: *1,25(OH)₂D₃*; *1,3-O-Diacetyl-1,25 Dihydroxy-16-ene-*

24-Keto-19-nor Cholecalciferol (compound (2)); and the compounds Mycophenolate Mofetil (MMF) and Dexamethasone (DEX).

Immunoglobulin-like transcript-3 upregulation was evaluated by flow cytometric analysis. Briefly, cells were preincubated with 200 mg/ml human IgG (Sigma Chemical, St. Louis, Missouri) and subsequently stained with anti-human ILT3 antibody (see, e.g., Cella, M. et al. (1997) J. Exp. Med. 185:1743) followed by anti-mouse IgG-phycoerythrin (Jackson). Cells were then analyzed with a FACScan[®] flow cytometer using a Cell Quest[®] software program (both from Beckton Dickinson, Mountain View, California).

10

II. Results

Incubation of monocyte-derived immature dendritic cells with 1,25(OH)₂D₃ (i), or 1,3-O-Diacetyl-1,25 Dihydroxy-16-ene-24-Keto-19-nor Cholecalciferol (compound (2)) upregulated the expression of ILT3 on their cell surface (see, Figure 1 below).

15 These data show that these vitamin D compounds upregulate ILT3 expression. Conversely, MMF and DEX, agents that also target dendritic cells, fail to upregulate ILT3 expression at any concentration tested, up to 1000 nM.

20




Figure 1

20

EXAMPLE 5

Soft Gelatin Capsule Formulation I

Item	Ingredients	mg/Capsule
1	Compound (2) from Example 1	10.001-0.02
2	Butylated Hydroxytoluene (BHT)	0.016
25	3 Butylated Hydroxyanisole (BHA)	0.016
4	Miglyol 812 qs.	160.0

Manufacturing Procedure:

1. BHT and BHA is suspended in Miglyol 812 and warmed to about 50 °C with stirring, until dissolved.
2. 1,25-dihydroxy-16-ene-24-oxo-5,6-trans-calcitriol is dissolved in the solution from step 1 at 50 °C.
3. The solution from Step 2 is cooled at room temperature.
4. The solution from Step 3 is filled into soft gelatin capsules.

Note: All manufacturing steps are performed under a nitrogen atmosphere and protected from light.

EXAMPLE 6

5

Soft Gelatin Capsule Formulation II

Item	Ingredients	mg/Capsule
1	Compound (2) from Example 1	10.001-0.02
2	di- α -Tocopherol	0.016
10 3	Miglyol 812 qs.	160.0

Manufacturing Procedure:

1. Di- α -Tocopherol is suspended in Miglyol 812 and warmed to about 50 °C with stirring, until dissolved.
- 15 2. 1,25-dihydroxy-16-ene-24-oxo-5,6-trans-calcitriol is dissolved in the solution from step 1 at 50 °C.
3. The solution from Step 2 is cooled at room temperature.
4. The solution from Step 3 is filled into soft gelatin capsules.

20 Incorporation by Reference

The contents of all references (including literature references, issued patents, published patent applications, and co-pending patent applications) cited throughout this application are hereby expressly incorporated herein in their entireties by reference.

25


Equivalents

Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents of the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the

30 following claims.

CLAIMS

1. A vitamin D₃ compound of formula I:



5

wherein:

X₁ and X₂ are each independently H₂ or =CH₂, provided X₁ and X₂ are not both =CH₂;

- R₁ and R₂ are each independently, hydroxyl, OC(O)C₁-C₄ alkyl, OC(O)hydroxyalkyl, OC(O)haloalkyl, provided that R₁ and R₂ are not both hydroxyl;

R₃ and R₄ are each independently hydrogen, C₁-C₄ alkyl, hydroxyalkyl, or haloalkyl, or R₃ and R₄ taken together with C₂₀ form C₃-C₆ cycloalkyl; and


R₅ and R₆ are each independently C₁-C₄ alkyl, hydroxyalkyl, or haloalkyl; and pharmaceutically acceptable esters, salts, and prodrugs thereof.

15

2. The compound of claim 1, wherein X₁ and X₂ are each H₂.
3. The compound of claim 1, wherein R₃ is hydrogen and R₄ is C₁-C₄ alkyl.
4. The compound of claim 3, wherein R₄ is methyl.
5. The compound of claim 1, wherein R₅ and R₆ are each independently methyl, ethyl or fluoromethyl.

20

6. The compound of claim 1, wherein R_5 and R_6 are each methyl.
7. The compound of claim 1, wherein R_1 and R_2 are each independently
5 hydroxyl or $OC(O)C_1-C_4$ alkyl.
8. The compound of claim 1, wherein R_1 and R_2 are each $OC(O)C_1-C_4$ alkyl.
9. The compound of claim 1, wherein R_1 and R_2 are each acetyloxy.
- 10
10. The compound of claim 9, wherein said compound is:




15

11. A method for treating a subject for a vitamin D₃ associated state, comprising administering to said subject an effective amount of a vitamin D₃ compound, such that said subject is treated for said vitamin D₃ associated state, wherein said compound is:

20

5



wherein:

10

X_1 and X_2 are each independently H_2 or $=CH_2$, provided X_1 and X_2 are not both $=CH_2$;

R_1 and R_2 are each independently, hydroxyl, $OC(O)C_1-C_4$ alkyl, $OC(O)$ hydroxyalkyl, $OC(O)$ haloalkyl, provided that R_1 and R_2 are not both hydroxyl;

15

R_3 and R_4 are each independently hydrogen, C_1-C_4 alkyl, hydroxyalkyl, or haloalkyl, or R_3 and R_4 taken together with C_{20} form C_3-C_6 cycloalkyl; and

R_5 and R_6 are each independently C_1-C_4 alkyl, hydroxyalkyl, or haloalkyl; and pharmaceutically acceptable esters, salts, and prodrugs thereof.

20

12. The method of claim 11, wherein X_1 and X_2 are each H_2 .

13.

The method of claim 11, wherein R_3 is hydrogen and R_4 is C_1-C_4 alkyl.

14. The method of claim 13, wherein R_4 is methyl.

15. The method of claim 11, wherein R_5 and R_6 are each independently methyl, ethyl or fluoromethyl.

5

16. The method of claim 11, wherein R_5 and R_6 are each methyl.


17. The method of claim 11, wherein R_1 and R_2 are each independently hydroxyl or $OC(O)C_1-C_4$ alkyl.

10

18. The method of claim 11, wherein R_1 and R_2 are each $OC(O)C_1-C_4$ alkyl.

19. The method of claim 11, wherein R_1 and R_3 are each acetyloxy.

15 20. The method of claim 19, wherein said compound is:



20 21. The method of claim 11, wherein said vitamin D_3 associated state is an ILT3-associated disorder.

22. The method of claim 21, wherein said ILT3-associated disorder is an immune disorder.

23. The method of claim 22, wherein said immune disorder is an autoimmune disorder.

5 24. The method of claim 23, wherein said autoimmune disorder is selected from the group consisting of type 1 insulin-dependent diabetes mellitus, adult respiratory distress syndrome, inflammatory bowel disease, dermatitis, meningitis, thrombotic thrombocytopenic purpura, Sjogren's syndrome, encephalitis, uveitis, uveoretinitis, leukocyte adhesion deficiency, rheumatoid arthritis, rheumatic fever,
10 Reiter's syndrome, psoriatic arthritis, progressive systemic sclerosis, primary biliary cirrhosis, pemphigus, pemphigoid, necrotizing vasculitis, myasthenia gravis, multiple sclerosis, lupus erythematosus, polymyositis, sarcoidosis, granulomatosis, vasculitis, pernicious anemia, CNS inflammatory disorder, antigen-antibody complex mediated diseases, autoimmune haemolytic anemia, Hashimoto's thyroiditis, Graves disease,
15 habitual spontaneous abortions, Reynard's syndrome, glomerulonephritis, dermatomyositis, chronic active hepatitis, celiac disease, autoimmune complications of AIDS, atrophic gastritis, ankylosing spondylitis and Addison's disease.

25. The method of claim 22, wherein said immune disorder is transplant
20 rejection.

26. The method of claim 11, wherein said vitamin D₃ associated state is a disorder characterized by an aberrant activity of a vitamin D₃-responsive cell.

25 27. The method of claim 26, wherein said disorder comprises an aberrant activity of a hyperproliferative skin cell.

28. The method of claim 27, wherein said disorder is selected from psoriasis, basal cell carcinoma and keratosis.

30

29. The method of claim 26, wherein said disorder comprises an aberrant activity of an endocrine cell.

30. The method of claim 29, wherein said endocrine cell is a parathyroid cell and the aberrant activity is processing and/or secretion of parathyroid hormone.
31. The method of claim 30, wherein said disorder is secondary
5 hyperparathyroidism.
32. The method of claim 26, wherein said disorder comprises an aberrant activity of a bone cell.
- 10 33. The method of claim 32, wherein said disorder is selected from osteoporosis, osteodystrophy, senile osteoporosis, osteomalacia, rickets, osteitis fibrosa cystica, and renal osteodystrophy.
34. The method of claim 26, wherein said disorder is cirrhosis or chronic
15 renal disease.
35. The method of claim 11, wherein said subject is a mammal.
36. The method of claim 35, wherein said subject is a human.
20
37. The method of claim 11, wherein said vitamin D₃ compound is administered in combination with a pharmaceutically acceptable carrier.
38. A pharmaceutical composition, comprising an effective amount of a
25 vitamin D₃ compound and a pharmaceutically acceptable carrier, wherein said vitamin D₃ compound is a compound as claimed in any one of claims 1-10.
39. The pharmaceutical composition of claim 38, wherein said effective amount is effective to treat a vitamin D₃ associated state.
30
40. The pharmaceutical composition of claim 39, wherein said vitamin D₃ associated state is an ILT3-associated disorder.

41. The pharmaceutical composition of claim 39, wherein said vitamin D₃ associated state is a disorder characterized by an aberrant activity of a vitamin D₃-responsive cell.

5 42. The pharmaceutical composition of claim 41, wherein said disorder comprises an aberrant activity of a hyperproliferative skin cell.

43. The pharmaceutical composition of claim 42, wherein said disorder is selected from psoriasis, basal cell carcinoma and keratosis.

10

44. The pharmaceutical composition of claim 41, wherein said disorder comprises an aberrant activity of an endocrine cell.

45. The pharmaceutical composition of claim 41, wherein said endocrine cell
15 is a parathyroid cell and the aberrant activity is processing and/or secretion of parathyroid hormone.

46. The pharmaceutical composition of claim 45, wherein said disorder is secondary hyperparathyroidism.

20

47. The pharmaceutical composition of claim 41, wherein said disorder comprises an aberrant activity of a bone cell.

48. The pharmaceutical composition of claim 47, wherein said disorder is
25 selected from osteoporosis, osteodystrophy, senile osteoporosis, osteomalacia, rickets, osteitis fibrosa cystica, and renal osteodystrophy.

49. The pharmaceutical composition of claim 41, wherein said disorder is cirrhosis or chronic renal disease.

30

50. The pharmaceutical composition of claim 41, wherein said disorder is in a mammalian subject.

51. The pharmaceutical composition of claim 501, wherein said subject is a human.
52. A method of ameliorating a deregulation of calcium and phosphate metabolism, comprising administering to a subject a therapeutically effective amount of a compound of any one of claims 1 to 10, so as to ameliorate the deregulation of the calcium and phosphate metabolism.
53. The method of claim 52, wherein the deregulation of the calcium and phosphate metabolism leads to osteoporosis.
54. A method of modulating the expression of an immunoglobulin-like transcript 3 (ILT3) surface molecule in a cell, comprising contacting said cell with a compound of any one of claims 1-10 in an amount effective to modulate the expression of an immunoglobulin-like transcript 3 (ILT3) surface molecule in said cell.
55. The method of claim 54, wherein said cell is within a subject.
56. A method of treating an ILT3-associated disorder in a subject, comprising administering to said subject a compound of any one of claims 1-10 in an amount effective to modulate the expression of an ILT3 surface molecule, thereby treating said ILT3-associated disorder in said subject.
57. The method of claim 56, wherein said ILT3-associated disorder is an immune disorder.
58. The method of claim 57, wherein said immune disorder is an autoimmune disorder.
59. The method of claim 58, wherein said autoimmune disorder is selected from the group consisting of type 1 insulin-dependent diabetes mellitus, adult respiratory distress syndrome, inflammatory bowel disease, dermatitis, meningitis, thrombotic thrombocytopenic purpura, Sjogren's syndrome, encephalitis, uveitis,

uveoretinitis, leukocyte adhesion deficiency, rheumatoid arthritis, rheumatic fever, Reiter's syndrome, psoriatic arthritis, progressive systemic sclerosis, primary biliary cirrhosis, pemphigus, pemphigoid, necrotizing vasculitis, myasthenia gravis, multiple sclerosis, lupus erythematosus, polymyositis, sarcoidosis, granulomatosis, vasculitis, pernicious anemia, CNS inflammatory disorder, antigen-antibody complex mediated diseases, autoimmune haemolytic anemia, Hashimoto's thyroiditis, Graves disease, habitual spontaneous abortions, Reynard's syndrome, glomerulonephritis, dermatomyositis, chronic active hepatitis, celiac disease, autoimmune complications of AIDS, atrophic gastritis, ankylosing spondylitis and Addison's disease.

10

60. The method of claim 57, wherein said immune disorder is transplant rejection.

61. A method of inducing immunological tolerance in a subject, comprising administering to said subject a compound of any one of claims 1-10 in an amount effective to modulate the expression of an ILT3 surface molecule, thereby inducing immunological tolerance in said subject.

15

62. The method of claim 61, wherein said immunological tolerance is induced in an antigen-presenting cell.

20

63. The method of claim 62, wherein said antigen-presenting cell is selected from the group consisting of dendritic cells, monocytes, and macrophages.

64. A method of inhibiting transplant rejection in a subject comprising administering to said subject a compound of any one of claims 1-10 in an amount effective to modulate the expression of an ILT3 surface molecule, thereby inhibiting transplant rejection in said subject.

25

65. The method of claim 64, wherein said transplant is a solid organ transplant.

30

66. The method of claim 64, wherein said transplant is a pancreatic islet transplant.

67. The method of claim 64, wherein said transplant is a bone marrow
5 transplant.

68. The method of any one of claims 52, 55, 56, 61 or 64, wherein said vitamin D₃ compound is administered to the subject using a pharmaceutically-acceptable formulation.

10

69. The method of claim 68, wherein said pharmaceutically-acceptable formulation provides sustained delivery of said vitamin D₃ compound to a subject for at least four weeks after the pharmaceutically-acceptable formulation is administered to the subject.

15

70. A packaged formulation for use in the treatment of an ILT-3 associated disorder, comprising a pharmaceutical composition comprising a compound of any one of claims 1-10 and a pharmaceutically-acceptable carrier packaged with instructions for use in the treatment of an ILT3-associated disorder.

20

71. A method for modulating immunosuppressive activity by an antigen-presenting cell, comprising contacting an antigen-presenting cell with a compound of any one of claims 1-9 in an amount effective to modulate ILT3 surface molecule expression, thereby modulating said immunosuppressive activity by said antigen-
25 presenting cell.

72. The method of any one of claims 52, 55, 56, 61 or 64 wherein said subject is a mammal.

30 73. The method of claim 72, wherein said subject is a human.

74. The method of claim 54, wherein said cell is an antigen-presenting cell.

75. The method of claims 71 or 74, wherein said antigen-presenting cell is selected from the group consisting of dendritic cells, monocytes, and macrophages.

76. The method of any one of claims 54, 56, 61 or 64, wherein the expression
5 of said immunoglobulin-like transcript 3 (ILT3) surface molecule is upregulated.

77. The method of any one of claims 11, 52, 54, 56, 61 or 64, wherein said compound is administered orally.

10 78. The method of any one of claims 11, 52, 54, 56, 61 or 64, wherein said compound is administered intravenously.

79. The method of any one of claims 11, 52, 54, 56, 61 or 64, wherein said compound is administered topically

15

80. The method of any one of claims 11, 52, 54, 56, 61 or 64, wherein compound is administered parenterally.

81. The method of any one of claims 54, 56, 61 or 64, wherein said
20 compound is administered at a concentration of 0.001 µg – 100 µg/kg of body weight.

82. The method of claim 81, where said compound is 1,3-O-Diacetyl-1,25-dihydroxy-16-ene-24-keto-19-nor-cholecalciferol

25 83. A packaged formulation for use in the treatment of a vitamin D₃ associated state, comprising a pharmaceutical composition comprising a compound of any one of claims 1-10 and instructions for use in the treatment of a vitamin D₃ associated state.

30 85. The method of claim 26, wherein the disorder comprises aberrant activity of a cell that expresses renin.

86. The method of claim 86, wherein the disorder is hypertension

87. The method of claims 86, wherein the compound of formula I suppresses expression of renin, thereby treating the subject for hypertension.
- 5 88. The method of claim 26, wherein the disorder is benign prostate hypertrophy.
89. The method of claim 26, wherein the disorder is neoplastic disease.
- 10 90. The method of claim 89, wherein the cancer is selected from the group consisting of leukemia, lymphoma, melanoma, osteosarcoma, colon cancer, rectal cancer, prostate cancer, and malignant tumors of the lung, breast, gastrointestinal tract, genitourinary tract.
- 15 91. The method of claim 26, wherein the disorder is neuronal loss.
92. The method of claim 91, wherein the disorder is selected from the group consisting of Alzheimer's Disease, Pick's Disease, Parkinson's Disease, Vascular Disease, Huntington's Disease, and Age-Associated Memory Impairment.
- 20 93. The method of claim 26, wherein the disorder is characterized by an aberrant activity of a vitamin D₃-responsive smooth muscle cell.
94. The method of claim 93, wherein the disorder is hyperproliferative
- 25 vascular disease selected from the group consisting of hypertension-induced vascular remodeling, vascular restenosis, and atherosclerosis.
95. The method of claim 93, wherein the disorder is characterized by an aberrant metabolism of a vitamin D₃-responsive smooth muscle cell.
- 30 96. The method of claim 95, wherein the disorder is arterial hypertension.
97. The compound of claim 1, wherein the haloalkyl is fluoroalkyl.

98. The compound of claim 97, wherein the fluoroalkyl is fluoromethyl or trifluoromethyl.
- 5 99. The method of claim 11, wherein the haloalkyl is fluoroalkyl.
100. The method of claim 99, wherein the fluoroalkyl is fluoromethyl or trifluoromethyl.
- 10 101. A method for treating a subject for a urogenital disorder, comprising administering to said subject an effective amount of a vitamin D₃ compound of formula I of claim 1, such that said subject is treated for said urogenital disorder.
102. The method of claim 101, wherein said disorder is bladder dysfunction.
- 15 103. The method of claim 102, wherein said bladder dysfunction is characterized by the presence of bladder hypertrophy.
104. The method of claim 101, wherein said disorder is interstitial cystitis.
- 20 105. The method of claim 104, wherein said interstitial cystitis is characterized by the presence of symptoms of bladder dysfunction and bladder inflammation.
106. The method of claim 101, wherein the disorder is benign prostatic
- 25 hyperplasia.



Generaldirektion 1

Directorate General 1

Direction générale 1

Teuten, Andrew John
Sagittarius Intellectual
Property Consultants Ltd
Taylor House
39 High Street
Marlow, Bucks. SL7 1AF
GRANDE BRETAGNE



EPO Customer Services

Tel.: +31 (0)70 340 45 00

Date

17.11.06


Reference BXL-P095/EPw	Application No./Patent No. 04789014.0 - 2103 PCT/US2004031412
Applicant/Proprietor Bioxell S.p.a.	

COMMUNICATION

The European Patent Office herewith transmits as an enclosure the supplementary European search report under Article 157(2)(a) EPC for the above-mentioned European patent application.

If applicable, copies of the documents cited in the European search report are attached.

- ☐ Additional set(s) of copies of the documents cited in the European search report is (are) enclosed as well.



Refund of the search fee

If applicable under Article 10 Rules relating to fees, a separate communication from the Receiving Section on the refund of the search fee will be sent later.



ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.

EP 04 78 9014

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

07-11-2006

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2004098507 A2	18-11-2004	EP 1677802 A2	12-07-2006
US 5976784 A	02-11-1999	NONE	

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82



European Patent
Office

**SUPPLEMENTARY
EUROPEAN SEARCH REPORT**

Application Number
EP 04 78 9014

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (IPC)
E	WD 2004/098507 A2 (BIOXELL S P A [IT]; ADORINI LUCIANO [IT]; PENNA GIUSEPPE [IT]; USKOKOV) 18 November 2004 (2004-11-18) * the whole document *	1-56	INV. C07C401/00 A61K31/593 A61P37/00 A61P35/00
D, X	US 5 976 784 A (DELUCA HECTOR F [US] ET AL) 2 November 1999 (1999-11-02) * the whole document *	1-56	
			TECHNICAL FIELDS SEARCHED (IPC)
			C07C A61K A61P
The supplementary search report has been based on the last set of claims valid and available at the start of the search.			
Place of search Munich		Date of completion of the search 7 November 2006	Examiner HORNICH-PARAF, E
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X: particularly relevant if taken alone Y: particularly relevant if combined with another document of the same category A: technological background O: non-written disclosure P: intermediate document</p> <p>T: theory or principle underlying the invention E: earlier patent document, but published on, or after the filing date D: document cited in the application L: document cited for other reasons &: member of the same patent family, corresponding document</p>			

1
EPO FORM 1503 (03.02.06) (P02004)